

53

Public Health Reports

VOLUME 58 DECEMBER 31, 1943 NUMBER 53

IN THIS ISSUE

Surveys of Mental Disorder in Sample Populations



CONTENTS

	Page
A survey of statistical studies on the prevalence and incidence of mental disorder in sample populations. Paul Lemkau, Christopher Tietze, and Marcia Cooper.....	1909
Deaths during week ended December 18, 1943:	
Deaths in a group of large cities in the United States.....	1927
Death claims reported by insurance companies.....	1927
PREVALENCE OF DISEASE	
United States:	
Reports from States for week ended December 25, 1943, and comparison with former years.....	1928
Weekly reports from cities:	
City reports for week ended December 11, 1943.....	1933
Rates, by geographic divisions, for a group of selected cities....	1935
Territories and possessions—Hawaii Territory—Honolulu—Dengue fever.....	1935
Foreign reports:	
Canada—Provinces—Communicable diseases—Week ended November 27, 1943.....	1936
Sweden—Notifiable diseases—October 1943.....	1936
Switzerland—Notifiable diseases—April–June 1943.....	1937
World distribution of cholera, plague, smallpox, typhus fever, and yellow fever—	
Cholera.....	1937
Plague.....	1937
Smallpox.....	1938
Typhus fever.....	1939
Yellow fever.....	1940

Public Health Reports

Vol. 58 • DECEMBER 31, 1943 • No. 53

A SURVEY OF STATISTICAL STUDIES ON THE PREVALENCE AND INCIDENCE OF MENTAL DISORDER IN SAMPLE POPULATIONS¹

By PAUL LEMKAU, CHRISTOPHER TIETZE, and MARCIA COOPER, *The Johns Hopkins University School of Hygiene and Public Health, Baltimore, Md.*

Intelligent planning for action in the field of mental hygiene requires knowledge of the magnitude of the problem. It is therefore necessary to ascertain the prevalence and incidence of mental hygiene problems in the general population. "Prevalence" is understood as referring to the number of cases active at a given moment, "incidence" to the number of new cases occurring either during an arbitrary period, usually a year, or during the life span of a group of individuals. In the latter case the rates are often referred to as "expectancies."

Attempts at complete enumeration of mental deviants have been made at various times in many countries in connection with the national censuses. Such attempts have been generally unsuccessful because of widespread failure on the part of informants and enumerators to recognize or report any but the most obvious cases.

Machinery for the general reporting of new cases of mental disorder does not exist in any country and for obvious reasons is not likely to be established. It is believed that in communities with an adequate and well established system of mental hospitals most psychotics find their way into these hospitals at some time during their lives, even though they spend a considerable part of their period of illness on the outside. For such populations rates of first admissions have been used as a substitute for incidence rates (18, 28, 32, 41). This method however, is not suitable for forms of mental disorder which do not as a rule require or receive institutional care.

The difficulties standing in the way of even approaching complete coverage of all cases of mental disorder in any large geographical or political unit have thus far proved insurmountable. The only practicable way yet discovered to approach the problem is to gather

¹This study was made with the support of the International Health Division of the Rockefeller Foundation.

information about prevalence and incidence in population samples. Such investigations have been carried out in various parts of the world.

The present paper is a critical survey of these studies. We have endeavored to report on all major investigations conducted and published during the last 15 years. There are few earlier studies of this type and these do not furnish much information of value because psychiatric concepts have changed radically since they were made. The progressive breakdown of international scientific relations makes it somewhat uncertain that all investigations have been discovered in the literature. All studies included cover a wide range of mental hygiene problems and were made on reasonably "average" populations. Investigations of single specific problems or on highly selected populations have been omitted.

Attention will first be given to five studies of *prevalence*. The earliest of these surveys was made in 1930 by Carl Brugger (8) in 116 villages in Thuringia with a total population of about 38,000. Only four of these communities have over a thousand inhabitants. The area is mixed agricultural-industrial in character. Most of the people are Protestants.

In the following years Brugger made two more surveys, one in five villages of the Bavarian Allgäu (10) and one in six villages east of the town of Rosenheim (11). These surveys were part of a larger study intended to cover several areas with a total population of about 50,000 inhabitants. We have been unable to find any other publications on this investigation besides Brugger's two papers. The Allgäu and Rosenheim areas are very similar in character; both are predominantly agricultural and almost exclusively Catholic. As the same technique was used in both instances and the total population was less than 9,000 the two appear combined in this review under the name "Bavaria survey."

In 1935 Erik Strömngren (40) carried out a mental hygiene survey on an island in the Baltic Sea, Bornholm, which belongs politically to Denmark. It has a population of 46,000; the chief sources of income are farming, fishing, quarrying, and tourist trade. The results of Strömngren's investigation are available in book form, including analyses of both prevalence and incidence of mental disorder.

Turning now to the United States we find two large community surveys of mental hygiene problems, one in Baltimore and one in Tennessee.² The Baltimore survey is limited to the Eastern Health District, an area of about one square mile in the eastern part of the city. The District serves as the field laboratory of the Johns Hopkins University School of Hygiene and Public Health. Two surveys were made here,

² Both supported by the International Health Division of the Rockefeller Foundation.

one in 1933 (13, 14, 15, 16) and a second in 1936 (20, 21, 22, 23). All further statements in this paper refer to the survey of 1936. At this time the Eastern Health District had 55,000 inhabitants of whom 23 percent were Negroes. Among the whites there are many families of Hebrew and of Czech extraction. The area is for the most part residential, and the income level for both white and Negro families is definitely below the average for the city.

Finally there is the survey of Williamson County, Tennessee (34), giving information as of September 1, 1938. Williamson County is a fairly typical agricultural county in middle Tennessee with an area of 586 square miles and a population of 25,000. About 78 percent of the people are native whites, mostly of English and Scotch-Irish extraction; 22 percent are Negroes. There is only one incorporated town, Franklin, with about 4,000 inhabitants.

It will be noted that none of the surveyed populations is anything like a representative sample of the national population of which it is a part, be it Germany, Denmark, or the United States. Bornholm is the only one of the survey areas which might be called a "natural" unit. Future investigators in this field will probably spend greater efforts upon the proper selection of sample populations. A well coordinated survey, conducted simultaneously in several urban and rural areas of the United States or other large countries, should yield more complete information on the prevalence of mental disorder than has hitherto been available. The classic mental deficiency survey (27) made in England from 1925 to 1927 might well serve as a model.

Bona fide residents of the survey areas who have become inmates of institutions located elsewhere are included in all studies.

In the following analysis the term "case finding" is used for the basic procedures by which individuals presenting problems in the field of mental hygiene become known to the investigator. The determination of the specific nature of the problem, and of the degree of deviation, is called diagnosis. The two may be carried out either separately or as one operation.

Brugger initiated his Thuringia survey by questionnaires sent to physicians, clergymen, teachers, and mayors in the area. Additional material was obtained by oral inquiries from the relatives of patients already known and from "older inhabitants" of the several villages. The files of hospitals, institutions, and prisons were also searched, in one case as far back as 30 years. Almost all reported cases were personally seen and diagnosed by the author. He mentions that he also examined a large number of individuals described to him as normal and that he feels satisfied as to the good mental health of these persons.

The investigation in Bavaria was much more intensive. The population was prepared by publicity in the local newspapers, by

announcement from the pulpit by the parish priests, and by notices on the communal bulletin boards by the mayors. Subsequently every family in the area was visited by the psychiatrist. A fairly complete history was taken for each individual and the diagnosis made on the basis of a personal examination. Reports from hospitals, institutions, physicians, teachers, etc., were utilized in checking the information given by the family. In this study case finding and diagnosis were combined.

Procedures in the Bornholm survey were closely patterned after Brugger's investigation in Thuringia. The records of mental and general hospitals, almshouses, and nursing homes were searched and oral information obtained from physicians, aldermen, clergymen, teachers, city officials, and "older inhabitants." Most cases were visited and examined by the author.

It should be mentioned that Strömngren also carried out a very intensive survey of one village on the island with the help of the local medical practitioners. This community, however, had only 900 inhabitants and therefore this phase of his investigation is not reviewed here in detail. It may be worth mentioning that the general prevalence rate computed for this village is five to six times as high as that found for the whole island by the less inclusive general investigation. This is approximately the same ratio as between Brugger's surveys in Bavaria and Thuringia.

In the Baltimore survey there was no reporting of cases directly to the investigating group. Case finding was carried out entirely by perusal of the written records of various institutions and agencies. This procedure was made possible by the high development of health and social services in the area under study. The total number of sources was 43, including, among others, public and private mental hospitals, training schools for mental defectives and for delinquents, psychiatric clinics, social agencies, certain departments of the public school system, the juvenile court, police and criminal courts, and the National Health Survey. In many of the cases thus discovered a psychiatric diagnosis or a more or less complete description by a competent social worker was available. Others, however, were included on the basis of circumstantial evidence. Diagnosis, where not already given in the medical sources, was made from the written records after careful study of the data. No cases were personally examined by the staff psychiatrist. This is probably a serious shortcoming in the investigative technique.

The material for the Tennessee survey was partly reported by key persons in the community—physicians, nurses, teachers, ministers, judges, postmasters, country storekeepers, etc.—and partly discovered by the field workers of the study, who spent years in the town participating extensively in many community activities of various kinds.

Institutional records were also searched. Somewhat over half of the cases were interviewed or examined by one or more members of the staff which consisted of a psychiatrist, social workers, and nurses.

In addition to this general investigation, an intensive house-to-house survey was conducted in three selected areas. Details are not yet available for this phase of the study beyond the fact that the prevalence rate for all types of problems was about twice as high in these areas as in the remainder of the county.

Considerable differences exist between the five surveys as to the types of mental hygiene problems included and as to the grouping into diagnostic categories. The three European studies are limited to psychiatric conditions in the narrower sense.

The Thuringia survey covers all "psychoses"—this term as used by Brugger includes epilepsy and hysteria—and, in addition, neurasthenia, psychopathic personalities so badly maladjusted that they themselves or their families suffered severely, chronic alcoholics showing moral, social, or economic deterioration, some so-called eccentrics, a few "neurologic" patients, and, finally, imbeciles and idiots. That the cases of mental deficiency were diagnosed on the basis of clinical evidence rather than by standardized tests is evident from the statement that the mental age of the imbeciles is between 6 and 12 years and the mental age of the idiots less than 6 years. These mental ages obviously do not describe the same intelligence levels as they would in this country. So far as the major psychoses are concerned, German and American terminology appear to be reasonably comparable, at least as regards the more common diseases, such as schizophrenia, manic-depressive psychosis, general paresis, arteriosclerotic psychosis, etc.

Scope and classification in the Bavarian investigation are essentially the same except that a large group of morons ("debil") and retarded individuals is included.

Strömngren states that he enumerated only the "psychoses" in his survey of Bornholm, but from the list of diagnostic groupings it appears that the term "psychosis" again covers not only what we generally mean thereby but also epilepsy and conditions which would be classified as psychoneuroses or minor psychoses in American practice. Most of Strömngren's subgroups can be identified easily although some of the terms are unfamiliar, as for example "depressio mentis" (reactive depression), "confusio mentis" (delirium), and a few more. Mental defectives were not systematically searched for but the author believes that the majority of the more severe oligophrenics are included. These cases are described as requiring special care and would probably have been classified as idiots or imbeciles in this country.

The American surveys are more ambitious and cover a much wider range of mental hygiene problems. The Baltimore group uses the following system of classification (20) which is planned to designate all known maladjusted individuals.

- Psychosis.
- Psychoneurosis.
- Psychopathic personality.
- Personality disorder in adults:
 - Psychotic traits.
 - Neurotic traits.
 - Psychopathic traits.
 - Behavior deviation.
- Behavior disorder in children:
 - Neurotic traits.
 - Conduct problems.
- Minor or possible disorder in adults and children.
- Epilepsy.
- Mental deficiency (I. Q. less than 70).
- School progress problems without mental deficiency.
- Adult delinquency without other information.

Under the title "personality disorders in adults" four different groups of cases are combined. The first three of the groups may be termed subclinical—that is, they include individuals in whom the clinical picture was not complete enough to justify their placement in a more sharply defined category. The fourth group covers a wide array of severely maladjusted personalities not otherwise definable and chiefly characterized by their interpersonal difficulties. The maladjustment ranges all the way from aggressiveness to submissiveness, including traits like deception, irritability, quarrelsomeness, uncooperativeness, chronic dependency, inefficiency, shiftlessness, suggestibility, unreliability, and so on. Severe alcoholics who are not psychotic are also included here.

"Behavior disorder in children" has two subdivisions: neurotic traits and conduct disturbances. Widely different problems are covered by the term "neurotic traits," including tics, habit spasms, stuttering, temper tantrums, enuresis, nail biting, and so on. Conduct problems are chiefly juvenile delinquency and school misbehavior.

The group called "minor or possible disorder" is very heterogeneous. It contains some cases with adequate records but minor problems, and others with very scanty information suggesting serious maladjustment.

In the Tennessee survey a somewhat different system of classification (34) was used but it includes about the same range of mental hygiene problems as is covered in Baltimore. The only group of cases in the Tennessee tabulations which has no counterpart in Baltimore is a number of individuals who because of constitutional or environmental handicaps are considered liable to become mental

hygiene problems although they were still well enough adjusted when investigated. The material is presented in seven major groupings.

- Psychosis.
- Psychoneurosis.
- Conduct and behavior problems.
- Psychopathic traits.
- Special personality types.
- Mental deficiency.
- Organic and miscellaneous conditions.

The "conduct and behavior problems" embrace major and minor delinquency of adults, juvenile delinquency, alcoholism, marital maladjustment, sexual promiscuity, etc. "Psychopathic traits" are such adult personality types as the schizoid, cyclothymic, and hysterical, and also the so-called neurotic traits of childhood, ties, enuresis, temper tantrums, etc. The group of "special personality types" includes, in addition to various minor deviations, many persons of borderline intelligence. Among the "organic and miscellaneous" conditions we find the epileptics, patients suffering from various diseases of the brain and nervous system, persons with endocrine and other general diseases and congenital abnormalities, cripples, and individuals living in an unfavorable home environment. The large majority of cases in this group are listed only as potential mental hygiene problems.

By definition a study of prevalence is based upon the number of cases active on a given day which may be called the census day for the particular survey. For certain types of problems in the field of mental hygiene the determination of activity status presents great difficulties, both practical and theoretical. The situation is simple in conditions of a permanent nature like mental deficiency where a case known to be present on census day must also be active. It is less simple in psychoses and similar illnesses; by careful investigation, however, it is possible to determine the time of onset or recovery within reasonable limits. But how long is a person a "delinquent" before, after, or between his criminal acts? This conundrum can only be answered arbitrarily.

Brugger does not discuss the question of activity status in his papers, but it appears that he counts as psychotics all individuals who have been mentally disordered at any time during their lives. This attitude probably reflects the importance attached to the constitutional factor by the German school of psychiatry.

Strömngren presents separately the numbers of active and recovered cases only for his total series of "psychotics" which includes epileptics and many psychoneurotics. The distinction is not made for each diagnostic group separately.

The Baltimore investigation includes all cases active at some time

during the survey year, and no others with the exception that an effort was made to ascertain the number of residents who had previously suffered from a psychosis. Such individuals, however, are not counted as active. The prevalence rates of the Baltimore survey are one-year rather than one-day rates. They cannot be converted easily into true prevalence rates.

In Tennessee it was possible to follow up all reported cases and to verify not only their continued residence in the county on census day but also to determine their activity status at that time. Among the active cases a further distinction was made between personal problems affecting only the patient and his immediate family and social problems involving the community at large. In each of these categories, severe, moderate, and minor deviations are distinguished. Such tabulations are available for the seven major diagnostic groups.

The possibilities of reaction inherent in the human personality are so manifold that it is small wonder that many individuals presenting mental hygiene problems require double or multiple diagnoses. This becomes particularly apparent in the two American prevalence surveys with their broad conception of the field. In both instances the difficulty has been overcome by the establishment of a rank order of diagnostic groups, descending, by and large, from more severe to less severe deviations. Each case with multiple diagnoses is assigned to the highest ranking group represented. This is called the leading classification or primary diagnosis and is generally, but not always, the most important condition as far as the individual is concerned. In addition there is available for certain diagnostic groups the total number of cases falling into that group irrespective of primary diagnosis. Among these groups is mental deficiency which often exists coincidentally with other conditions.

Brugger's papers report only a few cases with double diagnoses and those are listed individually. The Bornholm survey includes three duplications but no details are available.

It cannot fall into the scope of this review to reproduce in detail the findings of each survey. For complete information the reader is referred to the original publications. It does seem proper, however, to present in summary form the numbers of cases found in each diagnostic group and the crude prevalence rates per 1,000 general population, using the author's own system of classification. Later—and with considerable hesitation—the attempt will be made to rearrange the material for purposes of comparison. In all tables the range of sampling variation is indicated by 95 percent confidence limits.

Table 1 summarizes the findings of Brugger's two surveys in Thuringia and Bavaria. Although the population covered in the latter study was only one-fourth of that in the former, the number of cases found is larger. The prevalence rate for the aggregate of all forms of

mental disorder is almost five times as high. This difference is partly due to the inclusion of morons and persons with borderline mental deficiency and partly to the fact that a personal investigation was made by the psychiatrist of every family included in the Bavaria survey. This is suggested by the fact that the discrepancy appears largest for those types of mental disorder which are most likely to escape recognition and reporting by lay persons.

TABLE 1.—Cases of mental disorder in the Thuringia and Bavaria surveys

[Population: Thuringia survey 37,561; Bavaria survey 8,628]

	Number of cases		Rate per 1,000		95 percent confidence limits	
	Thuringia	Bavaria	Thuringia	Bavaria	Thuringia	Bavaria
Schizophrenia	73	22	1.9	2.5	1.5-2.4	1.5-3.6
Manic-depressive	20	12	.5	1.4	.3-.8	.6-2.2
Epilepsy	26	12	.7	1.4	.4-1.0	.6-2.2
Cerebral arteriosclerosis	13	3	.3	.3	.2-.5	.0-.7
Hysteria	32	7	.9	.8	.6-1.1	.2-1.4
Infectious psychosis	11		.3		.1-.5	
Undiagnosed psychosis	16	7	.4	.8	.2-.6	.2-1.4
Neurasthenia	15	7	.4	.8	.2-.6	.2-1.4
Psychopathic personality	25	24	.7	2.8	.4-.9	1.7-3.9
Alcoholism	15	22	.4	2.6	.2-.6	1.5-3.6
Eccentrics	6	14	.2	1.6	.0-.3	.8-2.8
Imbeciles and Idiots	201	137	5.4	15.9	4.6-6.1	13.2-18.5
Morons		158		18.3		15.5-21.2
Retarded		88		10.2		8.1-12.3
Miscellaneous conditions ¹	36	12	1.0	1.4	.6-1.3	.6-2.2
Total	479	517	12.8	59.9	11.6-13.9	54.8-65.1
Duplications	10	8				

¹ General paresis, cerebral syphilis, senile and presenile dementia, climacteric and reactive depression, puerperal psychosis, postencephalitic states, brain tumor, sequelae to concussion of the brain, migraine, postoperative tetany, and exogenous oligophrenia. None of these groups had over 10 cases in either survey.

Strömberg's figures for Bornholm are presented in table 2. His material does not include psychopathic personalities, alcoholics, and eccentrics and is therefore smaller in scope than Brugger's; otherwise the picture seems to be reasonably similar.

TABLE 2.—Active and recovered cases of mental disorder in the Bornholm survey

[Population: 45,930]

	Number of cases	Rate per 1,000	95 percent confidence limits
Schizophrenia	150	3.3	2.7-3.8
Manic-depressive	122	2.7	2.2-3.1
Epilepsy	47	1.0	.7-1.3
Psychogenic psychoses	34	.7	.5-1.0
Hysterical psychoses	35	.8	.5-1.0
Paranoid psychosis	13	.3	.1-.6
Confusio mentis	15	.3	.2-.5
Depressio mentis	49	1.1	.8-1.4
Other psychoses ¹	22	.5	.3-.7
Atypical psychoses	15	.3	.1-.4
Undiagnosed psychoses	28	.6	.4-.8
Oligophrenia	191	4.2	3.6-4.7
Total	716	15.6	14.4-16.7
Duplications	3		

¹ General paresis; psychoses with organic brain disease; senile, presenile, and arteriosclerotic dementia; and alcoholic psychoses. There were less than 10 cases in each of these groups.

The data for the Eastern Health District of Baltimore shown in table 3 introduce a quite different type of coverage, reaching far beyond the ranks of individuals manifestly mentally disordered. The psychotics, psychoneurotics, and psychopathic personalities who may together be called the "clinical" group furnish only 17 percent of all cases and even the inclusion of the "subclinical" types does not raise this proportion to more than 20 percent. It should be noted that table 3 presents the cases active in 1936 by leading classification. A complete tabulation would reveal hundreds of cases with double

TABLE 3.—Active cases of mental disorder in the Baltimore survey

[Population: 55,129]

Leading classification	Number of cases	Rate per 1,000	95 percent confidence limits
Psychosis.....	367	6.7	6.0-7.3
Schizophrenia.....	158	2.9	2.4-3.3
Manic-depressive.....	41	.7	.5-1.0
Senile and arteriosclerotic.....	38	.7	.5-.9
Alcoholic.....	15	.3	.1-.4
Syphilitic.....	29	.5	.3-.7
With mental deficiency.....	28	.5	.3-.7
Other ¹	27	.5	.3-.7
Undiagnosed.....	31	.6	.4-.8
Psychoneurosis.....	171	3.1	2.6-3.6
Psychopathic personality.....	30	.5	.4-.7
Personality disorder in adults.....	218	4.0	3.4-4.5
Psychotic traits.....	26	.5	.3-.7
Neurotic traits.....	60	1.1	.8-1.4
Psychopathic traits.....	13	.2	.1-.4
Behavior deviation.....	119	2.2	1.8-2.5
Behavior disorder in children.....	449	8.1	7.4-8.9
Neurotic traits.....	162	2.9	2.5-3.4
Conduct problems.....	287	5.2	4.6-5.8
Minor and possible disorder in adults and children.....	651	11.8	10.9-12.7
Epilepsy.....	75	1.4	1.1-1.7
Mental deficiency.....	375	6.8	6.1-7.5
School progress problems without mental deficiency.....	434	7.9	7.1-8.6
Adult delinquency without other information.....	567	10.3	9.4-11.1
Total active cases ²	3,337	60.5	58.5-62.6

¹ Involuntal, with epilepsy, post traumatic, and deliria not due to alcohol.² Active+inactive cases: 3,416=62.0 per 1,000.

or multiple diagnosis, indicating a much more comprehensive analysis than was undertaken by Brugger and Strömngren. Some groups would expand spectacularly. The total number of epileptics, for instance, was 126 or 2.3 (1.9-2.7) per 1,000, that of mental defectives 694 or 12.6 (11.7-13.5) per 1,000 general population.

Table 4 summarizes the basic information about the active and inactive cases in the Tennessee survey. Again the full-fledged psychoses and psychoneuroses are very definitely in the minority. Mentally defective individuals not presenting a definite social or personal problem are listed as inactive here. The total number of mental defectives is 376 or 15.2 (13.6-16.7) per 1,000.

TABLE 4.—*Active and inactive cases of mental disorder in the Tennessee Survey*

[Population: 24,804]

Primary diagnosis	Number of cases			Rate per 1,000 (total cases)	95 percent confidence limits
	Active	Inactive	Total		
Psychosis.....	121	35	156	6.3	5.3- 7.3
Schizophrenia.....			43	1.7	1.2- 2.3
Affective.....			41	1.7	1.1- 2.2
Senile.....			23	.9	.5- 1.3
With mental deficiency.....			15	.6	.3- .9
Other ¹			24	1.0	.6- 1.4
Undiagnosed.....			10	.4	.2- .7
Psychoneurosis.....	89	10	99	4.0	3.2- 4.8
Conduct and behavior disorder.....	285	129	414	16.7	15.1-18.3
Psychopathic traits.....	152	34	186	7.5	6.4- 8.6
Special personality traits.....	208	127	335	13.5	12.1-15.0
Mental deficiency.....	19	184	203	8.2	7.1- 9.3
Organic and miscellaneous conditions.....	40	288	328	13.2	11.8-14.7
All types.....	914	807	1,721	69.4	66.1-72.7

¹ General paresis, other organic states, posttraumatic, with alcoholism, and with epilepsy.

Is there any legitimate possibility for comparison of findings between the five surveys reviewed in this paper? They were made by four different investigators (or groups of investigators) in three countries. The widely divergent methods and practices employed in each study for case finding and classification has been set forth in the preceding pages as well as the great variations in scope. Nevertheless there is a strong temptation to seek a basis for comparison. It is realized that such a comparison cannot be exact; that it cannot give more, in fact, than a very general idea nor be more than a very slim basis for generalizations. Apart from the inevitable sampling variation, the range of which is indicated in the tables, a difference of prevalence rates between two surveys may be due to differences in scope, in case finding, in diagnosis, or to actual variations in the true prevalence of the various types of mental disorder. At the present state of our knowledge no valid conclusions can yet be drawn from the differences between the observed prevalence rates. On the other hand if the rate is found reasonably near the same level in all surveys, then the suspicion seems justified that we may be on the track of a numerical relationship of wider applicability.

The prevalence rates computed from the total numbers of active and inactive cases included in each survey are 12.8 per 1,000 general population for Thuringia, 15.6 for Bornholm, 59.9 for Bavaria, 62.0 for Baltimore, and 69.4 for Tennessee. In view of what has been said before it is obvious that the marked differences are due mainly to variations in scope and that at least some of the apparent similarities of rates must be entirely coincidental.

A fairly good picture of the prevalence of the major forms of mental disorder may be obtained from a combination of the psychoses and psychoneuroses, covering what might be called the field of "clinical"

or "traditional" psychiatry. A clear-cut separation of the two groups is not feasible in the three European surveys. Likewise it is necessary to include both active and inactive cases as this distinction cannot be made in Brugger's papers. The aggregate prevalence rate is 6.4 (5.6-7.2) per 1,000 general population in Thuringia, 8.7 (6.7-10.7) in Bavaria, 10.3 (9.0-11.5) in Tennessee, 11.4 (10.5-12.4) in Bornholm, and 11.8 (10.9-12.7) in Baltimore.³ It should be borne in mind, however, that the comparability is by no means absolute, all epileptics, for instance, being included in Thuringia, Bavaria, and Bornholm, but only epileptics with psychosis in the American surveys. Nevertheless the statement is probably not far off the mark that about 1 percent of the general population is or has been suffering from major mental disorder, that is from a psychosis or psychoneurosis. Among these persons there seem to be considerably more psychotics than psychoneurotics. Such would, of course, not be the case if all individuals with any "neurotic" manifestations were included.

Table 5 presents a synopsis of prevalence rates for two supposedly uniformly defined groups. Active and inactive cases are included in the figures for the two "endogenous" psychoses and it will also be

TABLE 5.—Active and recovered cases of schizophrenia and manic-depressive psychosis in five surveys

	Schizophrenia	Manic-depressive	Schizophrenia and manic-depressive
Number of cases:			
Thuringia.....	73	20	93
Bavaria.....	22	12	34
Bornholm.....	150	122	272
Baltimore ¹	193	62	255
Tennessee.....	43	41	84
Rate per 1,000:			
Thuringia.....	1.9	.5	2.5
Bavaria.....	2.5	1.4	3.9
Bornholm.....	3.3	2.7	5.9
Baltimore ¹	3.5	1.1	4.6
Tennessee.....	1.7	1.7	3.4
95 percent confidence limits:			
Thuringia.....	1.5-2.4	.3-.8	2.0-3.0
Bavaria.....	1.5-3.6	.6-2.2	2.6-5.3
Bornholm.....	2.8-3.8	2.2-3.1	5.2-6.6
Baltimore ¹	3.0-4.0	.8-1.4	4.1-5.2
Tennessee.....	1.2-2.2	1.1-2.2	2.6-4.1

¹ One-day estimate made on the basis of the known numbers of hospitalized and nonhospitalized patients and of postpsychotic individuals in each diagnostic group.

noted that the aggregate rates present a more regular pattern than the constituent parts. No other groups appear comparable throughout all five surveys. Unfortunately the numbers of cases underlying many of the rates in table 5 are quite small.

Turning now to attempts at estimating the *incidence* of mental disorder for sample populations we find a number of papers using a

³ The Baltimore figure does not include inactive cases of psychoneurosis. In the Tennessee study there were only 10 such individuals, corresponding to a rate of 0.4 per 1,000. Figures in parentheses indicate 95 percent confidence limits. Apparent inconsistencies with the tables are due to multiple diagnoses.

method inaugurated by Professor Rüdin (33) of the German Research Institute for Psychiatry at the Kaiser Wilhelm Institute in Munich. Some of these studies were conducted and published under the auspices of the Institute; others were made independently but conform to the technique worked out there.

In each of them the procedure consists of three phases; first, construction of the sample population, second, case finding and diagnosis, and third, statistical analysis. The construction of the sample population starts with the selection of a group of *propositi*. These must meet a number of requirements which are often hard to reconcile. The group should be of adults and as large as can be handled, of average social-economic status, accessible to the psychiatrist but unbiased as far as disposition to mental disorder is concerned. Actually most of the groups of *propositi* studied number between 100 and 200 persons. They fall by type of origin into three classes; first, wives and husbands of patients with organic psychoses; second, medical and surgical patients themselves; and third, nonpatient material collected on a geographic or occupational basis.

The next step is to obtain a roster of all siblings of the *propositi*. In order to get complete information this must be done with great thoroughness and often the help of interested members in the family of the *propositus* is enlisted. Corroborative evidence is obtained from church and civil authorities, schools, etc. Psychiatric case finding is very closely connected with the genealogic work on the sample population. The informants are questioned about illnesses, commitments to mental institutions, outstanding personality traits, and other pertinent facts. Wherever possible the opinions of several observers about the same individual are checked against each other. The picture is rounded out by hospital records, physicians' statements, and other documents. Most of the authors have endeavored to see personally as many individuals in the families of the *propositi* and in particular as many of the mentally abnormal cases reported to them as feasible. In every study, however, some of the siblings are dead, some live in remote places, and a few are uncooperative. Therefore some of the psychiatric diagnoses must be made on the basis of descriptions by lay persons.

The German school has made conscious efforts to keep diagnostic practice and terminology as uniform as possible. Some groupings appear to be comparable to their counterparts in American psychiatry. This is true for schizophrenia, manic-depressive psychoses, general paresis, and idiopathic epilepsy. In some cases distinctions are somewhat unsatisfactory as, for instance, between senile dementia and cerebral arteriosclerosis and perhaps also between hysteria and psychopathic personality. The latter term apparently is used for a much wider range of conditions in Germany than in this country.

The greater number of the genealogic studies covers only institutionalized cases of psychopathic personality. Data on mental deficiency are likewise not directly comparable to American and English studies because our highly standardized testing procedures are not used and diagnoses are made on the basis of social history and clinical observation. The term "oligophrenia" covers idiots and imbeciles. For mental deficiency of a lesser degree the expression "debil" is used, but many of the papers do not attempt to report these cases. Of alcoholics, only such individuals are included who show evident moral, social, or economic decline or at least lack of progress which ordinarily would be expected (24). This criterion is applicable on all occupational and cultural levels.

Of prime importance for a successful estimate of the incidence of mental disorder from genealogical data is of course the cooperativeness of the *propositi*, their relatives, and other informants. It must be made clear to them why such information is sought; the disinclination to reveal abnormalities in the family has to be overcome and they must be convinced that the investigation will not lead to any discrimination against them. It is reported that after the enactment of the German sterilization laws people became quite reluctant to assist in genealogical studies of their own families (4, 6). We have been unable to find any papers of the type described published after 1937. It may well be that the work had to be discontinued because it was felt that the results had become too unreliable.

The final step is the statistical analysis. Incidence is given in terms of expectancy, expectancy being defined as the chance of developing or acquiring a specific disease if the individual lives through the age period of susceptibility. It is obvious that a rate thus defined is significant for conditions that have their onset at birth, in youth, or middle life, but that it cannot be used for instance for senile dementia because here the period of susceptibility has no upper age limit. The German group responsible for the method was and probably still is chiefly interested in the heredity of mental disorder and has not paid much attention to the psychoses of advanced age.

The actual computations of expectancy are usually carried out for the major psychoses by the abridged method of Weinberg (42). This consists in putting into the numerator the number of cases and into the denominator the total number of siblings who have passed through the period of susceptibility plus one-half of those who have entered but not passed it. The age of susceptibility is assumed to be 16 to 40 years for schizophrenia, 20 to 50 years for manic-depressive psychosis, and 30 to 50 years for general paresis. For mental deficiency, epilepsy, and other forms of mental disorder the number of cases is as a rule simply related to the number of siblings who have passed their tenth

birthday. These approximations seem close enough for most purposes. More complicated formulae, though sometimes used, appear unjustified in view of the scanty basic data.

In general there has been a marked tendency on the part of the German workers to overestimate the statistical validity of their findings. As early as 1928 when only two studies had been published, Luxenburger wrote as follows: "The expectancy of dementia praecox, manic-depressive psychoses, epilepsy, and general paresis seems to be *practically established* for a predominantly urban average population (Munich): this expectancy is for dementia praecox 0.85 percent, for manic-depressive psychosis 0.41 percent, for epilepsy 0.29 and for general paresis 1.73 percent. The figures for dementia praecox and paresis in particular are stable enough to be considered as *standard rates*" (24). The italics are Luxenburger's. These rates were carried from one publication to the other and often uncritically accepted as more or less official. It is not generally known that these so-called standard rates are based on eight cases of paresis, five of schizophrenia, and two each of manic-depressive psychosis and epilepsy. In subsequent years when one paper after the other appeared in the *Zeitschrift für die gesamte Neurologie und Psychiatrie* the authors could not always resist the temptation to compare their individual findings with those of others. Practically all these comparisons are worthless because of the small numbers in each sample. From the scientific point of view, it would appear that a better way of utilizing the genealogical method would be to make as many studies as possible with the same technique and to combine them into one large series. Such compilations have been made from time to time in Germany at least for certain diagnostic groups (12, 17, 37, 38).

Table 6 presents a synopsis based upon 15 papers published between 1927 and 1937. In four (7, 24, 31, 35) the *propositi* are wives and husbands of patients with organic psychoses, in six (2, 3, 5, 6, 36, 43) they are medical and surgical patients and in the remaining five (1, 4, 9, 19, 25) a nonpatient group has been used. Four (5, 24, 35, 36) groups of *propositi* were investigated in Munich, four (9, 19, 25, 43) in the rural part of Bavaria, two (4, 6) in Silesia, one each in Berlin (31) and Saxony (1), and three in Switzerland (2, 3, 7). Northern Germany is definitely under-represented. The total number of *propositi* in the 15 studies is 2,090 and the total number of siblings 10,684. Table 6 presents in summary form the number of cases of mental disorder found among these siblings and the expectancies calculated from them. Weinberg's abridged method is used for schizophrenia, manic-depressive psychosis, and general paresis; all other forms of illness are related to persons over 10 years of age, except alcoholism where those over 20 are used as a population base. The confidence limits indicate that the sample is not as numerically adequate as one

would like to have it, apart from the question of representativeness. The almost complete absence of senile dementia and cerebral arteriosclerosis—one case of either is included among the “other psychoses”—is of course due to the age distribution of the siblings. Only 15 percent of them are over 50 years of age and very few over 60.

TABLE 6.—*Mental disorder found among a total number of 10,884 siblings of 2,090 propiiti in 15 papers from Germany and Switzerland*

	Bibliography No.															Total	Rate per 1,000	95 percent confidence limits
	1	2	3	4	5	6	7	9	10	24	25	31	35	36	43			
Schizophrenia.....	7	6	2	3	4	1	6	3	---	3	2	1	2	2	---	42	7.7	5.4-10.1
Manic-depressive.....	3	1	---	1	---	1	---	1	---	1	---	1	1	---	---	10	2.3	.9-3.8
General paresis.....	2	---	---	---	1	---	---	---	---	---	4	1	---	---	---	12	3.5	1.5-5.5
Epilepsy.....	6	4	3	3	1	2	1	1	2	1	2	3	1	2	1	33	4.2	2.8-5.7
Hysteria.....	6	5	---	---	---	---	---	---	2	1	2	---	---	---	---	17	2.2	1.1-3.2
Other psychoses.....	---	---	---	---	1	1	---	2	---	---	---	---	8	---	---	7	.9	.2-1.6
Undiagnosed psychoses.....	---	---	---	1	1	---	1	1	---	2	---	---	---	1	---	7	.9	.2-1.6
Oligophrenia.....	9	14	5	12	6	6	3	6	5	1	1	2	2	5	2	76	9.7	7.5-11.9
Morons (“debil”).....	(1)	(1)	(1)	(1)	(1)	(1)	7	3	6	(1)	3	13	(1)	2	13	147	15.5	11.0-19.9
Psychopathic personalities:	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
Institutionalized.....	---	---	---	3	3	2	2	---	---	---	---	2	1	3	---	16	2.0	1.0-3.0
Not institutionalized.....	34	2	(1)	12	(1)	2	5	(1)	(1)	(1)	(1)	26	(1)	4	(1)	185	18.8	14.8-22.8
Alcoholism.....	1	20	6	---	4	---	4	2	5	5	3	1	6	3	---	60	8.6	6.4-10.8

¹ Morons (“debil”) and psychopathic personalities, not institutionalized, are included in only 7 papers each.

Luxenburger was quite lucky in the case of schizophrenia with his original “standard rate” of 0.85 percent based on five cases, but he was less successful concerning manic-depressive psychosis and epilepsy and quite off the mark in general paresis.

The combined incidence for all forms of psychosis and psychoneurosis—excluding the disorders of advanced age but including epilepsy—appears to be about 20 per 1,000 individuals living through the age period of susceptibility. This is reasonably compatible with the figures on prevalence previously presented.

Outside Germany and Switzerland, the genealogic method has been used by Eliot Slater (39) in England, who used a group of surgical patients as propiiti and considers his results as most unsatisfactory, and by Strömngren (40) in his investigation of Bornholm. Strömngren’s findings based on a material of 427 propiiti with 1,927 siblings are in line with the German figures, but the number of cases is of course quite small. No such studies have been made in the United States. Any undertaking of this type would be beset with great difficulties in a country where general registration of population is not practiced.

The expectancies based on genealogic studies conducted in Germany and Switzerland may be compared with estimates for New York State based on first admissions to mental hospitals (41). This com-

parison seems justified for schizophrenia, manic-depressive psychosis, and general paresis. Using a suitable life table the German figures can be converted to incidence rates in terms of persons born rather than of persons surviving the period of susceptibility. The reduction is carried out in table 7. The German rates are consistently much lower than the rates for New York State. This may be entirely due to incomplete case finding in the genealogical material but it is also possible that the incidence of the three psychoses is actually higher in New York. Several considerations point in this direction. Almost two-thirds of the population of New York State live in the metropolitan area of the largest city of the world; only five percent live on farms. The groups from which the German samples are drawn are much less urbanized and most psychoses seem to be more common in urban than in rural areas. It is also likely that the presence of three million foreign-born tends to elevate the incidence rate for New York. There is evidence that psychoses occur more often among migrants—both interstate and intercontinental—than in nonmigrant populations (26, 29, 30). On the other hand it must not be forgotten that the estimates for New York are based upon first admissions and do not include cases who at no time receive intramural care. This limitation does not apply to the genealogic sample studies.

TABLE 7.—*Expectancy per 1,000 born for three major psychoses in Germany and New York State*

	Germany	New York
Schizophrenia.....	6.8	16.0
Manic-depressive.....	2.0	5.0
General paresis.....	2.9	4.4

In summary, then, it appears that poor selection of sample populations and insufficient numbers of cases as well as differences in investigative methods, differences in fundamental concepts, and differences in diagnosis and classification tend to make the available studies of prevalence and incidence of mental disorder basically incomparable. In this sense the result of this "survey of surveys" is disappointing. At the same time it furnishes a challenge to continue investigation and to make effort to correct the shortcomings of the science in which we work.

REFERENCES

- (1) Berlitz, B.: Beitrag zur Belastungsstatistik einer Durchschnittsbevölkerung. *Zeitschr. f. d. ges. Neurol. u. Psychiat.*, **152**: 622-643 (1935).
- (2) Bleuler, M.: Psychotische Belastung von körperlich Kranken. *Zeitschr. f. d. ges. Neurol. u. Psychiat.*, **142**: 780-810 (1932).
- (3) Bleuler, M., and Rapoport, L.: Untersuchungen über die konstitutionelle Verwandtschaft von Tuberkulose und Geisteskrankheiten. *Zeitschr. f. d. ges. Neurol. u. Psychiat.*, **153**: 649-679 (1935).

- (4) Boeters, D.: Belastungsstatistik einer schlesischen Durchschnittsbevölkerung. *Zeitschr. f. d. ges. Neurol. u. Psychiat.*, **155**: 675-701 (1936).
- (5) Boeters, H.: Familienuntersuchungen bei einer Durchschnittsbevölkerung. *Zeitschr. f. d. ges. Neurol. u. Psychiat.*, **153**: 90-116 (1935).
- (6) Bormann, H.: Zur Belastungsstatistik der Durchschnittsbevölkerung. *Zeitschr. f. d. ges. Neurol. u. Psychiat.*, **159**: 110-135 (1937).
- (7) Brugger, C.: Zur Frage einer Belastungsstatistik der Durchschnittsbevölkerung. *Zeitschr. f. d. ges. Neurol. u. Psychiat.*, **118**: 458-488 (1929).
- (8) Brugger, C.: Versuch einer Geisteskrankenzählung in Thüringen. *Zeitschr. f. d. ges. Neurol. u. Psychiat.*, **133**: 352-390 (1931).
- (9) Brugger, C.: Psychiatrisch-genealogische Untersuchungen an einer Allgäuer Landbevölkerung. *Zeitschr. f. d. ges. Neurol. u. Psychiat.*, **145**: 516-544 (1933).
- (10) Brugger, C.: Psychiatrische Ergebnisse einer medizinischen, anthropologischen und soziologischen Bevölkerungsuntersuchung. *Zeitschr. f. d. ges. Neurol. u. Psychiat.*, **147**: 489-524 (1933).
- (11) Brugger, C.: Psychiatrische Bestandesaufnahme im Gebiete eines medizinisch-anthropologischen Zensus in der Nähe von Rosenheim. *Zeitschr. f. d. ges. Neurol. u. Psychiat.*, **160**: 189-201 (1938).
- (12) Brugger, C.: Übersicht über auslesefreie Untersuchungen in der Verwandtschaft Schwachsinniger. *Zeitschr. f. psych. Hyg.*, **11**: 97-120 (1938).
- (13) Cohen, B. M., and Fairbank, R. E.: Statistical contributions from the mental hygiene study of the Eastern Health District in Baltimore. I. General account of the 1933 mental hygiene survey of the Eastern Health District. *Am. J. Psychiat.*, **94**: 1153-1161 (1938).
- (14) Cohen, B. M., and Fairbank, R. E.: Statistical contributions from the mental hygiene study of the Eastern Health District in Baltimore. II. Psychosis in the Eastern Health District. *Am. J. Psychiat.*, **94**: 1377-1395 (1938).
- (15) Cohen, B. M., Fairbank, R. E., and Greene, E.: Statistical contributions from the mental hygiene study of the Eastern Health District in Baltimore. III. Personality disorder in the Eastern Health District in 1933. *Human Biology*, **11**: 112-129 (1939).
- (16) Cohen, B. M., Tietze, C., and Greene, E.: Statistical contributions from the mental hygiene study of the Eastern Health District in Baltimore. IV. Further studies on personality disorder in the Eastern Health District in 1933. *Human Biology*, **11**: 485-512 (1939).
- (17) Conrad, K.: Übersicht über auslesefreie Untersuchungen in der Verwandtschaft von Epileptikern. *Zeitschr. f. psych. Hyg.*, **10**: 167-184 (1937).
- (18) Dorn, H. F.: The incidence and future expectancy of mental disease. *Pub. Health Rep.*, **53**: 1991-2004 (1938).
- (19) Göppel, W.: Untersuchung der näheren Verwandtschaft von Allgäuer Reichsbahnangestellten. *Zeitschr. f. d. ges. Neurol. u. Psychiat.*, **113**: 445-478 (1928).
- (20) Lemkau, P., Tietze, C., and Cooper, M.: Mental hygiene problems in an urban district. *Mental Hygiene*, **25**: 624-646 (1941).
- (21) Lemkau, P., Tietze, C., and Cooper, M.: Mental hygiene problems in an urban district. Second paper. *Mental Hygiene*, **26**: 100-119 (1942).
- (22) Lemkau, P., Tietze, C., and Cooper, M.: Mental hygiene problems in an urban district. Third paper. *Mental Hygiene*, **26**: 275-288 (1942).
- (23) Lemkau, P., Tietze, C., and Cooper, M.: Mental hygiene problems in an urban district. Fourth paper. *Mental Hygiene*, **27**: 279-295 (1943).
- (24) Luxenburger, H.: Demographische und psychiatrische Untersuchungen in der engeren biologischen Familie von Paralytikerehegatten. *Zeitschr. f. d. ges. Neurol. u. Psychiat.*, **112**: 331-491 (1928).
- (25) Magg, F.: Zur Belastungsstatistik der Durchschnittsbevölkerung. *Zeitschr. f. d. ges. Neurol. u. Psychiat.*, **119**: 39-68 (1929).
- (26) Malzberg, B.: Migration and mental disease among Negroes in New York State. *Am. J. Phys. Anthropol.*, **21**: 107-113 (1936).
- (27) Mental Deficiency Committee, being a Joint Committee of the Board of Education and the Board of Control: Report of the . . . Part IV. Report on an investigation into the incidence of mental deficiency in six areas, 1925-27, by E. O. Lewis. London, 1929.
- (28) Metropolitan Life Insurance Company: The chances of becoming mentally ill. *Statistical Bulletin*, **18** (7): 5-8 (1937).

- (29) Ødegaard, Ø.: Emigration and Insanity: A study of mental disease among the Norwegianborn population of Minnesota. Copenhagen, 1932. (*Acta psychiatrica et neurologica, supplementum IV.*)
- (30) Ødegaard, Ø.: Emigration and mental health. *Mental Hygiene*, **20**: 546-553 (1936).
- (31) Panse, F.: Beitrag zur Belastungsstatistik der Durchschnittsbevölkerung. *Zeitschr. f. d. ges. Neurol. u. Psychiat.*, **154**: 194-222 (1936).
- (32) Pollock, H. M., and Malzberg, B.: Expectation of mental disease. *Psychiat. Quart.*, **2**: 549-579 (1928).
- (33) Rüdin, E.: Zur Vererbung und Neuentstehung der Dementia praecox. Berlin, 1916.
- (34) Roth, W. F., and Luton, F. H.: The mental health program in Tennessee. *Am. J. Psychiat.*, **99**: 662-675 (1943).
- (35) Schulz, B.: Zur Frage einer Belastungsstatistik der Durchschnittsbevölkerung. *Zeitschr. f. d. ges. Neurol. u. Psychiat.*, **109**: 15-48 (1927).
- (36) Schulz, B.: Zur Belastungsstatistik der Durchschnittsbevölkerung. *Zeitschr. f. d. ges. Neurol. u. Psychiat.*, **136**: 386-411 (1931).
- (37) Schulz, B.: Übersicht über auslesefreie Untersuchungen in der Verwandtschaft Schizophrener. *Zeitschr. f. psych. Hyg.*, **9**: 130-156 (1936).
- (38) Schulz, B.: Übersicht über auslesefreie Untersuchungen in der Verwandtschaft Manisch-Depressiver. *Zeitschr. f. psych. Hyg.*, **10**: 39-60 (1937).
- (39) Slater, E.: The incidence of mental disorder. *Annals of Eugenics*, **6**: 172-185 (1935).
- (40) Strömberg, E.: Beiträge zur psychiatrischen Erblehre. Copenhagen, 1938. (*Acta psychiatrica et neurologica, supplementum XIX.*)
- (41) Tietze, C.: A note on the incidence of mental disease in the State of New York. *Am. J. Psychiat.*, **100**: 402-405 (November 1943).
- (42) Weinberg, W.: Zur Korrektur des Einflusses der Lebensdauer und Todesauslese auf die Ergebnisse bestimmter Kreuzungen. *Arch. f. Rassen- u. Gesellschaftsbiologie*, **11**: 434-444 (1915).
- (43) Wolf, G.: Untersuchung der näheren Verwandtschaft von Allgäuer Kropfoperierten. *Zeitschr. f. d. Ges. Neurolog. u. Psychiat.*, **117**: 728-752 (1928).

DEATHS DURING WEEK ENDED DECEMBER 18, 1943

[From the Weekly Mortality Index, issued by the Bureau of the Census, Department of Commerce]

	Week ended Dec. 18, 1943	Correspond- ing week, 1942
Data from 88 large cities of the United States:		
Total deaths.....	11,379	9,449
Average for 3 prior years.....	8,927	
Total deaths, 50 weeks of year.....	451,285	419,868
Deaths under 1 year of age.....	657	672
Average for 3 prior years.....	570	
Deaths under 1 year of age, 50 weeks of year.....	32,004	29,140
Data from industrial insurance companies:		
Policies in force.....	66,117,272	65,272,092
Number of death claims.....	12,220	12,006
Death claims per 1,000 policies in force, annual rate.....	9.6	9.6
Death claims per 1,000 policies, 50 weeks of year, annual rate.....	9.6	9.1

PREVALENCE OF DISEASE

No health department, State or local, can effectively prevent or control disease without knowledge of when, where, and under what conditions cases are occurring

UNITED STATES

REPORT FROM STATES FOR WEEK ENDED DECEMBER 25, 1943

Summary

A total of 83,973 cases of influenza was reported for the week, as compared with 82,951 for the preceding week. However, omitting Kentucky, which reported 34,148 cases for the week ended December 18, stated to be based on estimates in some localities, the figures for reported cases for the 2 weeks are 48,803 and 81,753, respectively. These figures probably provide a more nearly accurate index to the trend. In 44 States (excluding Kentucky and 3 New England States which reported no cases) increases were recorded currently in 32 States, decreases in 11 States, and the same number reported in 1 State. Reported cases for these States for weeks ended November 27, and December 4, 11, 18, and 25 are as follows: 2,464, 4,486, 18,330, 48,801, and 81,753. For the corresponding weeks of 1940 the figures for all States reporting were as follows: 1,332, 3,014, 9,663, 29,864, and 42,457. In the mild epidemic of that season, the peak was reached during the week ended January 18, 1941, with 107,270 reported cases.

An index to the corresponding increase in mortality is given by the following figures:

	Week ended—				
	Nov. 27	Dec. 4	Dec. 11	Dec. 18	Dec. 25
Deaths from influenza and pneumonia in 39 scattered cities:					
1943.....	254	381	450	832	1,063
1942.....	299	294	332	378	397
3-year average.....	284	292	304	334	338
Total deaths in 90 large cities.....	8,677	9,846	10,373	11,524	12,646
Percentage increase.....	-3.3	13.5	5.4	11.1	9.7

For the current week, the number of cases of poliomyelitis declined from 89 to 39. The 5-year median is 48. This is the first week since April that the incidence has been below the median.

A total of 361 cases of meningococcus meningitis was reported, as compared with 281 last week and a 5-year (1938-42) median of 34. The incidence increased in all of the nine geographic areas except the South Atlantic and Mountain. A total of 2,932 cases has been reported since the beginning of the fourth quarter of the year, as compared with 916 for the same period last year, the highest comparable incidence of the past 6 years. A total of 17,459 cases has been reported to date this year.

A total of 475 cases of dysentery (amebic, bacillary, and unspecified) was reported. Cumulative totals for 51 weeks of the year and for the past 12 weeks, respectively (comparable figures for last year in parentheses), are 24,396 (19,467) and 5,650 (3,506).

New low rates will apparently be established this year for both smallpox and typhoid fever. To date, 730 cases of smallpox and 5,418 cases of typhoid fever have been reported, as compared with 801 and 6,652 cases, respectively, last year, in which year both diseases established new low incidence rates.

A total of 4,475 cases of endemic typhus fever has been reported to date, as compared with 3,662 for the same period last year, the year of highest reported incidence.

The total deaths, all causes, in 90 large cities in the United States up to and including the week ended December 25 is 469,290, as compared with 433,740 for the same period last year.

Telegraphic morbidity reports from State health officers for the week ended December 25, 1943, and comparison with corresponding week of 1942 and 5-year median

In these tables a zero indicates a definite report, while leaders imply that, although none was reported, cases may have occurred.

Division and State	Diphtheria			Influenza			Measles			Meningitis, men- ingococcus		
	Week ended—		Med- ian 1938- 42	Week ended—		Med- ian 1938- 42	Week ended—		Med- ian 1938- 42	Week ended—		Med- ian 1938- 42
	Dec. 25, 1943	Dec. 26, 1942		Dec. 25, 1943	Dec. 26, 1942		Dec. 25, 1943	Dec. 26, 1942		Dec. 25, 1943	Dec. 26, 1942	
NEW ENGLAND												
Maine.....	1	1	1	62	1	1	91	13	37	4	5	1
New Hampshire.....	0	1	0	(1)	-----	-----	2	14	4	1	0	0
Vermont.....	0	0	0	-----	-----	-----	1	159	25	1	0	0
Massachusetts.....	7	4	4	-----	-----	-----	340	352	196	17	6	2
Rhode Island.....	0	0	0	35	-----	-----	86	1	1	6	0	0
Connecticut.....	1	1	0	832	2	2	6	211	67	7	2	0
MIDDLE ATLANTIC												
New York.....	14	19	20	2 475	10	14	653	243	395	43	6	3
New Jersey.....	6	6	7	351	12	8	473	38	38	24	4	0
Pennsylvania.....	10	7	17	63	3	-----	455	909	678	34	6	5
E. NO. CENTRAL												
Ohio.....	13	8	16	6,986	7	8	1,294	46	42	12	1	1
Indiana.....	7	8	8	677	9	14	85	75	11	22	8	0
Illinois.....	4	8	27	437	7	14	147	46	46	28	0	0
Michigan.....	6	11	8	304	1	2	539	45	206	15	1	0
Wisconsin.....	8	3	0	1,832	31	31	455	164	164	12	3	0
W. NO. CENTRAL												
Minnesota.....	7	2	2	16	1	2	442	2	31	2	1	0
Iowa.....	2	1	3	11,463	-----	3	33	64	69	1	0	0
Missouri.....	0	4	10	100	3	5	35	6	6	24	1	1
North Dakota.....	3	2	2	1,443	24	24	230	0	12	2	0	0
South Dakota.....	3	0	4	39	-----	-----	9	147	7	0	0	0
Nebraska.....	1	3	2	51	5	1	3	87	8	0	0	0
Kansas.....	5	8	5	975	7	15	25	25	70	5	1	1
SOUTH ATLANTIC												
Delaware.....	0	0	0	-----	-----	-----	12	0	1	0	0	0
Maryland.....	3	8	8	696	11	5	36	3	3	10	8	0
District of Colum- bia.....	0	0	1	845	3	3	25	0	3	2	1	1
Virginia.....	5	12	14	7,584	383	152	93	12	41	6	10	0
West Virginia.....	3	4	9	3,747	16	18	83	6	6	1	3	3
North Carolina.....	9	4	38	76	2	10	90	3	145	1	2	1
South Carolina.....	7	3	4	1,058	294	315	41	3	3	1	1	1
Georgia.....	5	6	10	1,405	71	71	68	13	18	4	2	0
Florida.....	8	0	4	114	1	9	10	1	2	5	0	1
E. SO. CENTRAL												
Kentucky.....	2	3	5	2,220	18	18	9	58	12	7	0	0
Tennessee.....	7	6	11	982	56	52	51	13	29	5	0	1
Alabama.....	9	15	15	1,573	143	143	56	1	20	5	1	1
Mississippi.....	0	5	5	-----	-----	-----	-----	-----	-----	2	0	1
W. SO. CENTRAL												
Arkansas.....	16	6	7	4,090	41	97	35	58	28	0	0	0
Louisiana.....	10	9	9	148	9	9	2	44	11	6	1	1
Oklahoma.....	2	8	8	2,022	94	97	9	103	11	4	0	0
Texas.....	20	21	47	9,392	823	823	50	16	35	4	2	2
MOUNTAIN												
Montana.....	0	0	0	2,654	15	15	126	26	26	4	0	0
Idaho.....	0	4	2	12	1	1	0	60	3	0	0	0
Wyoming.....	2	0	1	814	66	15	12	10	10	0	0	0
Colorado.....	8	8	11	1,041	34	36	116	27	27	0	3	3
New Mexico.....	0	0	1	28	3	2	4	3	10	1	0	0
Arizona.....	3	0	2	731	83	126	8	1	3	0	0	0
Utah.....	0	0	0	5,723	43	43	10	261	19	1	1	0
Nevada.....	0	0	0	908	-----	-----	0	5	0	0	0	0
PACIFIC												
Washington.....	2	0	1	3,200	4	3	18	311	182	5	1	1
Oregon.....	2	2	2	2,201	13	13	62	289	37	3	11	0
California.....	12	20	20	3,668	30	102	87	44	190	24	0	0
Total.....	234	241	424	83,973	2,290	2,693	6,532	4,018	4,544	361	92	34
51 weeks.....	13,489	15,236	16,569	293,567	105,727	182,255	594,435	500,085	500,085	17,459	3,587	1,986

See footnotes at end of table.

Telegraphic morbidity reports from State health officers for the week ended December 26, 1943, and comparison with corresponding week of 1942 and 5-year median—Con.

Division and State	Poliomyelitis			Scarlet fever			Smallpox			Typhoid and paratyphoid fever ¹		
	Week ended—		Median 1938-42	Week ended—		Median 1938-42	Week ended—		Median 1938-42	Week ended—		Median 1938-42
	Dec. 25, 1943	Dec. 26, 1942		Dec. 25, 1943	Dec. 26, 1942		Dec. 25, 1943	Dec. 26, 1942		Dec. 25, 1943	Dec. 26, 1942	
NEW ENGLAND												
Maine.....	0	0	0	22	8	16	0	0	0	1	0	1
New Hampshire.....	0	0	0	2	15	9	0	0	0	0	0	0
Vermont.....	0	0	0	4	1	7	0	0	0	0	0	0
Massachusetts.....	3	0	0	210	238	145	0	0	0	0	1	1
Rhode Island.....	0	0	0	9	3	3	0	0	0	0	0	0
Connecticut.....	0	0	0	30	28	33	0	0	0	2	0	0
MIDDLE ATLANTIC												
New York.....	1	4	1	265	279	297	0	0	0	1	3	6
New Jersey.....	0	0	0	79	47	92	0	0	0	2	0	0
Pennsylvania.....	1	0	1	163	157	245	0	14	0	0	0	7
EAST NORTH CENTRAL												
Ohio.....	1	0	1	220	243	243	0	13	1	1	0	3
Indiana.....	0	0	0	76	68	84	0	6	5	1	3	1
Illinois.....	2	0	1	136	168	323	3	1	1	0	4	4
Michigan ²	0	0	0	86	95	182	0	0	1	4	0	2
Wisconsin.....	0	0	1	141	204	153	0	0	1	0	0	0
WEST NORTH CENTRAL												
Minnesota.....	0	1	1	96	56	76	0	0	15	0	3	0
Iowa.....	0	1	1	97	48	70	0	0	2	0	0	1
Missouri.....	0	0	0	46	60	79	0	0	2	0	5	4
North Dakota.....	0	1	0	6	14	13	0	0	0	1	0	0
South Dakota.....	0	0	0	19	23	17	0	2	2	0	0	0
Nebraska.....	0	0	0	36	20	20	0	0	1	0	0	0
Kansas.....	0	0	0	60	46	85	0	0	0	0	2	0
SOUTH ATLANTIC												
Delaware.....	0	0	0	3	2	18	0	0	0	1	0	0
Maryland ³	0	1	1	47	40	40	0	0	0	1	1	2
District of Columbia.....	0	0	0	16	12	10	0	0	0	0	0	1
Virginia.....	0	1	1	40	45	31	0	0	0	1	3	3
West Virginia.....	0	0	1	38	37	67	0	0	0	0	0	1
North Carolina.....	1	0	0	36	39	68	0	0	0	0	0	0
South Carolina.....	0	0	0	5	11	10	0	0	0	0	2	1
Georgia.....	0	0	0	13	35	23	0	0	0	0	0	6
Florida.....	0	0	0	6	5	8	1	0	0	4	0	0
EAST SOUTH CENTRAL												
Kentucky.....	0	1	0	43	22	59	0	1	0	4	1	2
Tennessee.....	0	2	0	38	58	58	0	0	0	1	1	2
Alabama.....	0	0	1	13	22	25	1	0	0	0	3	2
Mississippi ³	0	2	0	10	2	8	0	0	0	0	1	1
WEST SOUTH CENTRAL												
Arkansas.....	0	1	1	4	4	12	0	0	0	0	1	3
Louisiana.....	1	0	1	6	4	11	0	0	0	3	4	4
Oklahoma.....	1	0	0	30	27	24	0	0	2	5	1	1
Texas.....	1	7	3	36	39	48	1	0	2	3	4	7
MOUNTAIN												
Montana.....	0	0	0	39	8	30	0	0	0	0	0	0
Idaho.....	0	1	0	7	4	7	0	0	0	0	0	0
Wyoming.....	0	0	0	4	46	11	0	0	0	0	0	0
Colorado.....	0	1	0	35	58	29	0	0	1	0	1	0
New Mexico.....	0	0	0	2	2	7	0	0	0	1	2	3
Arizona.....	1	2	0	5	0	4	0	0	0	0	2	1
Utah ³	6	0	0	52	54	13	0	0	0	0	0	0
Nevada.....	2	0	0	2	3	0	0	0	0	0	0	0
PACIFIC												
Washington.....	2	0	0	74	9	44	0	0	0	2	0	0
Oregon.....	6	0	0	79	11	11	0	0	0	2	0	0
California.....	10	10	3	196	103	107	0	0	0	1	1	3
Total.....	39	36	48	2,712	2,527	2,979	6	37	47	42	49	89
51 weeks.....	12,358	4,143	7,261	137,454	123,995	152,425	730	801	2,401	5,418	6,652	9,505

See footnotes at end of table.

Telegraphic morbidity reports from State health officers for the week ended December 25, 1943, and comparison with corresponding week of 1942 and 5-year median—Con.

Division and State	Whooping cough			Week ended Dec. 25, 1943									
	Week ended—		Median 1933-42	Anthrax	Dysentery			Encephalitis, infectious	Leptosy	Rocky Mt. spotted fever	Tularemia	Typhus fever	
	Dec. 25, 1943	Dec. 26, 1942			Amebic	Bacillary	Un-specified						
NEW ENGLAND													
Maine.....	6	39	19	0	0	0	0	0	0	0	0	0	
New Hampshire.....	0	5	7	0	0	0	0	0	0	0	0	0	
Vermont.....	10	42	35	0	0	0	0	0	0	0	0	0	
Massachusetts.....	54	194	194	0	0	4	0	2	0	0	0	0	
Rhode Island.....	7	26	26	0	0	0	0	0	0	0	0	0	
Connecticut.....	12	29	54	0	0	2	0	0	0	0	0	0	
MIDDLE ATLANTIC													
New York.....	125	321	410	0	2	2	0	4	1	0	0	0	
New Jersey.....	61	130	162	0	3	0	0	1	0	0	1	0	
Pennsylvania.....	66	253	253	0	0	0	0	0	0	0	0	0	
E. NO. CENTRAL													
Ohio.....	38	133	133	0	0	0	0	0	0	0	2	0	
Indiana.....	23	24	17	0	0	0	0	0	0	0	1	0	
Illinois.....	54	80	171	0	4	3	0	0	0	0	3	0	
Michigan ¹	75	191	203	0	0	1	0	1	0	0	0	0	
Wisconsin.....	94	142	142	0	0	0	0	0	0	0	0	0	
W. NO. CENTRAL													
Minnesota.....	12	30	42	0	4	0	0	0	0	0	0	0	
Iowa.....	18	28	22	0	0	0	0	0	0	0	0	0	
Missouri.....	4	11	20	0	0	0	0	0	0	0	0	0	
North Dakota.....	31	7	4	0	0	0	0	0	0	0	0	0	
South Dakota.....	0	1	2	0	0	0	0	0	0	0	0	0	
Nebraska.....	3	1	2	0	0	0	0	0	0	0	0	0	
Kansas.....	15	46	39	0	0	1	0	0	0	0	0	0	
SOUTH ATLANTIC													
Delaware.....	2	7	4	0	0	0	0	0	0	0	0	0	
Maryland ²	28	69	49	0	0	0	1	0	0	0	3	0	
District of Columbia.....	3	9	13	0	0	0	0	0	0	0	0	0	
Virginia.....	60	59	59	0	0	0	27	0	0	0	2	0	
West Virginia.....	13	4	15	0	0	0	0	0	0	0	0	0	
North Carolina.....	48	26	85	0	0	0	0	0	0	0	0	3	
South Carolina.....	41	4	19	0	0	3	0	0	0	0	0	1	
Georgia.....	0	13	10	0	0	2	0	0	0	0	0	16	
Florida.....	20	5	5	0	0	5	0	0	0	0	0	9	
E. SO. CENTRAL													
Kentucky.....	47	19	24	0	0	1	0	0	0	0	2	0	
Tennessee.....	19	16	19	0	1	0	0	0	0	0	0	0	
Alabama.....	12	43	36	0	0	0	0	0	0	1	0	12	
Mississippi ³				0	0	0	0	0	0	0	0	2	
W. SO. CENTRAL													
Arkansas.....	22	26	26	0	0	1	0	0	0	0	0	1	
Louisiana.....	1	0	5	0	0	0	0	0	0	0	0	4	
Oklahoma.....	0	15	5	0	0	0	0	0	0	0	0	0	
Texas.....	151	128	121	0	10	375	0	0	0	0	0	34	
MOUNTAIN													
Montana.....	6	17	6	0	0	0	0	1	0	0	0	0	
Idaho.....	1	1	4	0	0	0	0	0	0	0	0	0	
Wyoming.....	0	6	3	0	0	0	0	1	0	0	0	0	
Colorado.....	22	6	32	0	0	0	0	0	0	0	0	0	
New Mexico.....	8	9	15	0	0	0	0	0	0	0	0	0	
Arizona.....	20	0	10	0	0	0	14	0	0	0	1	0	
Utah ⁴	8	14	18	0	0	0	0	0	0	0	0	9	
Nevada.....	0	0	0	0	0	0	0	0	0	0	0	0	
PACIFIC													
Washington.....	20	16	16	0	0	0	1	0	0	0	0	0	
Oregon.....	18	9	12	0	0	0	0	0	0	0	0	0	
California.....	42	102	137	0	2	4	0	1	0	0	0	0	
Total.....	1,320	2,455	3,176	0	26	406	43	11	1	1	15	82	
51 weeks.....	175,128	175,284	175,284	65	2,104	17,968	4,324	680	30	436	780	4,475	
51 weeks, 1942.....				78	1,160	11,953	6,354	552	45	452	879	3,662	

¹ Upper respiratory infections, 21.

² New York City only.

³ Period ended earlier than Saturday.

⁴ Including paratyphoid fever cases reported separately as follows: Michigan, 1; New York, 1; Florida, 2.

⁵ Later reports for Kentucky reveal that an estimated 30,000 of the more than 35,000 cases occurring in the week ended Dec. 11 (Public Health Reports, Dec. 24, 1943, p. 1901, footnote 3) were included in the telegraphic report for the week ended Dec. 18.

WEEKLY REPORTS FROM CITIES

City reports for week ended Dec. 11, 1943

This table lists the reports from 88 cities of more than 10,000 population distributed throughout the United States, and represents a cross section of the current urban incidence of the diseases included in the table.

	Diphtheria cases	Encephalitis, infectious, cases	Influenza		Measles cases	Meningitis, meningococcus, cases	Pneumonia deaths	Pollomyelitis cases	Scarlet fever cases	Smallpox cases	Typhoid and paratyphoid fever cases	Whooping cough cases
			Cases	Deaths								
NEW ENGLAND												
Maine:												
Portland	0	0		0	11	0	2	0	7	0	0	4
New Hampshire:												
Concord	0	0		0	0	0	0	0	1	0	0	0
Vermont:												
Barre	0	0		0	0	0	0	0	0	0	0	0
Massachusetts:												
Boston	4	0		0	14	10	18	1	53	0	0	25
Fall River	0	0		0	1	0	0	1	3	0	0	0
Springfield	0	0		0	17	2	3	0	7	0	0	2
Worcester	0	0		0	1	0	9	0	61	0	0	7
Rhode Island:												
Providence	0	0	1	0	65	3	4	0	3	0	0	19
Connecticut:												
Bridgeport	0	0	1	1	0	1	4	0	4	0	0	0
Hartford	0	0	1	0	0	1	1	0	9	0	0	2
New Haven	0	0	1	0	2	1	2	0	2	0	0	1
MIDDLE ATLANTIC												
New York:												
Buffalo	0	0		2	3	3	11	1	9	0	0	2
New York	13	1	70	5	444	20	89	7	185	0	4	50
Rochester	1	0		2	0	2	26	0	5	0	0	10
Syracuse	0	0		0	0	0	3	0	3	0	0	12
New Jersey:												
Camden	2	0	1	1	0	1	3	0	3	0	0	1
Newark	0	0	12	0	1	3	8	0	9	0	0	11
Trenton	0	0	1	0	2	0	3	0	11	0	0	3
Pennsylvania:												
Philadelphia	2	0	13	5	5	15	44	0	41	0	1	12
Pittsburgh	1	0	9	5	155	1	23	0	20	0	2	6
Reading	0	0		0	1	0	3	0	1	0	0	5
EAST NORTH CENTRAL												
Ohio:												
Cincinnati	3	0	2	2	5	0	5	0	23	0	0	2
Cleveland	0	0	7	1	48	4	6	0	47	0	2	13
Columbus	0	0	1	1	18	0	3	0	12	0	0	1
Indiana:												
Fort Wayne	0	0		0	0	0	1	0	2	0	0	0
Indianapolis	6	0		3	2	2	18	0	18	0	0	4
South Bend	0	0		0	48	0	0	0	0	0	0	0
Terre Haute	0	0		0	0	0	3	0	0	0	0	0
Illinois:												
Chicago	2	0	120	8	8	13	42	4	67	0	1	23
Springfield	0	0	17	0	2	0	7	0	1	0	0	0
Michigan:												
Detroit	3	0	44	0	13	11	19	0	52	0	3	21
Flint	0	0		0	9	0	8	0	4	0	0	15
Grand Rapids	0	0		0	51	0	2	0	6	0	0	2
Wisconsin:												
Kenosha	0	0		0	1	0	0	0	5	0	0	1
Milwaukee	1	0	4	4	7	4	12	0	43	0	0	20
Racine	0	0		0	1	0	2	0	8	0	0	9
Superior	0	0		0	121	1	2	0	0	0	0	0
WEST NORTH CENTRAL												
Minnesota:												
Duluth	0	0		0	5	0	2	0	13	0	0	16
Minneapolis	7	0		2	56	2	5	0	50	0	0	5
Missouri:												
Kansas City	2	0	38	0	1	2	5	0	18	0	1	1
St. Joseph	0	0		0	0	0	0	0	3	0	0	0
St. Louis	0	0	40	0	5	6	12	1	14	0	0	14

City reports for week ended Dec. 11, 1943—Continued

	Diphtheria cases	Encephalitis, infectious cases	Influenza		Measles cases	Meningitis, meningococcus, cases	Pneumonia deaths	Pollomyelitis cases	Scarlet fever cases	Smallpox cases	Typhoid and paratyphoid fever cases	Whooping cough cases
			Cases	Deaths								
WEST NORTH CENTRAL—continued												
North Dakota:												
Fargo.....	0	0	—	0	15	0	0	0	2	0	0	
Nebraska:												
Omaha.....	2	0	—	0	1	0	2	0	15	0	0	0
Kansas:												
Topeka.....	1	0	—	0	1	0	0	0	4	0	0	6
Wichita.....	0	0	5	2	0	1	9	0	8	0	0	0
SOUTH ATLANTIC												
Delaware:												
Wilmington.....	0	0	—	0	2	0	0	0	4	0	0	0
Maryland:												
Baltimore.....	1	0	32	3	14	3	25	0	30	0	0	23
Cumberland.....	0	0	—	0	0	0	0	0	0	0	0	0
Frederick.....	0	0	—	0	0	0	0	0	0	0	0	0
District of Columbia:												
Washington.....	3	0	45	4	28	1	17	0	23	0	0	6
Virginia:												
Lynchburg.....	0	0	—	0	242	0	1	0	1	0	0	2
Richmond.....	0	0	13	1	10	3	1	0	1	0	0	3
Roanoke.....	0	0	—	0	1	0	0	0	1	0	0	2
West Virginia:												
Charleston.....	0	0	—	0	5	0	0	0	0	0	0	0
Wheeling.....	0	0	—	0	0	0	3	0	2	0	0	2
North Carolina:												
Raleigh.....	0	0	—	0	0	0	0	0	0	0	0	0
Winston-Salem.....	0	0	—	0	65	0	0	0	3	0	0	0
South Carolina:												
Charleston.....	0	0	32	1	1	2	2	0	2	0	0	0
Georgia:												
Atlanta.....	0	0	96	0	4	1	6	1	2	0	0	1
Brunswick.....	0	0	—	0	16	0	3	0	0	0	0	0
Savannah.....	0	0	4	0	0	1	1	0	1	0	0	0
Florida:												
Tampa.....	0	0	—	0	0	0	3	0	0	0	0	0
EAST SOUTH CENTRAL												
Tennessee:												
Memphis.....	0	0	44	3	1	2	8	0	6	0	0	2
Nashville.....	0	0	—	0	0	0	5	0	6	0	0	4
Alabama:												
Birmingham.....	1	0	17	1	4	0	1	0	0	0	0	0
Mobile.....	0	0	—	4	0	1	5	0	0	0	0	0
WEST SOUTH CENTRAL												
Arkansas:												
Little Rock.....	0	0	—	0	1	0	3	0	0	0	0	0
Louisiana:												
New Orleans.....	4	0	18	5	3	0	11	1	2	0	0	0
Shreveport.....	0	0	—	0	0	0	2	0	2	0	0	0
Texas:												
Dallas.....	0	0	1	1	0	1	4	1	1	0	0	6
Galveston.....	0	0	—	0	0	0	0	0	0	0	0	0
Houston.....	4	0	—	1	1	2	7	0	2	0	0	0
San Antonio.....	0	0	1	0	0	0	3	0	0	0	0	0
MOUNTAIN												
Montana:												
Billings.....	0	0	—	0	0	0	1	0	0	0	0	0
Great Falls.....	0	0	—	0	47	1	0	0	3	0	0	0
Helena.....	0	0	—	0	0	0	0	0	5	0	0	0
Missoula.....	0	0	—	0	0	0	3	0	1	0	0	0
Idaho:												
Boise.....	0	0	—	0	0	0	0	0	1	0	1	0
Colorado:												
Denver.....	0	0	55	2	11	0	10	0	13	0	0	12
Pueblo.....	0	0	—	0	130	0	2	0	1	0	0	11
Utah:												
Salt Lake City.....	0	0	—	1	4	1	1	1	22	0	0	1

City reports for week ended Dec. 11, 1943—Continued

	Diphtheria cases	Encephalitis, infectious, cases	Influenza		Measles cases	Meningitis, meningococcus, cases	Pneumonia deaths	Poliomyelitis cases	Scarlet fever cases	Smallpox cases	Typhoid and paratyphoid fever cases	Whooping cough cases
			Cases	Deaths								
PACIFIC												
Washington:												
Seattle.....	2	0	-----	1	4	1	4	0	4	0	0	6
Spokane.....	0	0	1	1	22	0	1	0	23	0	0	4
Takoma.....	2	0	0	0	3	0	0	0	11	0	0	3
California:												
Los Angeles.....	4	0	44	2	5	2	3	2	37	0	0	9
Sacramento.....	0	0	-----	0	3	1	1	1	2	0	0	0
San Francisco.....	0	0	1	0	2	4	13	3	12	0	0	2
Total.....	71	1	792	75	1,769	136	580	25	1,072	0	15	452
Corresponding week, 1942.....	87	0	152	53	1,073	35	440	11	847	0	8	1,048
Average, 1938-42.....	104	-----	663	132	982	-----	401	-----	877	6	21	1,137

Dysentery, amebic.—Cases: New York, 1; Philadelphia, 1; Detroit, 2; Los Angeles, 1; San Francisco, 1.

Dysentery, bacillary.—Cases: Bridgeport, 1; New York, 5; Rochester, 1; Syracuse, 1; Detroit, 3; Charleston, S. C., 4; Los Angeles, 8.

Dysentery, unspecified.—Cases: San Antonio, 6.

Leprosy.—Cases: Los Angeles, 1.

Tularemia.—Cases: Reading, 1; Chicago, 1.

Typhus fever.—Cases: Winston-Salem, 1; Atlanta, 1; Brunswick, 1; Savannah, 3; Little Rock, 1; New Orleans, 1; Houston, 1; San Antonio, 15 (10 delayed reports are included); Los Angeles, 2.

1 3-year average, 1940-42.

2 5-year median.

Rates (annual basis) per 100,000 population, by geographic groups, for the 88 cities in the preceding table (estimated population, 1942, 34,443,000)

	Diphtheria case rates	Encephalitis, infectious, case rates	Influenza		Measles case rates	Meningitis, meningococcus, case rates	Pneumonia death rates	Pollomyelitis case rates	Scarlet fever case rates	Smallpox case rates	Typhoid and paratyphoid fever case rates	Whooping cough case rates
			Case rates	Death rates								
New England.....	9.9	0.0	9.9	2.5	276	44.7	106.8	5.0	373	0.0	0.0	149
Middle Atlantic.....	2.5	0.4	47.3	8.0	272	20.1	95.0	3.6	128	0.0	3.1	54
East North Central.....	2.8	6.0	113.9	11.1	195	20.4	75.9	2.3	166	0.0	2.5	76
West North Central.....	26.2	0.0	181.5	8.7	184	24.1	76.5	2.2	278	0.0	2.2	92
South Atlantic.....	6.8	0.0	379.4	15.4	663	18.6	121.3	1.7	120	0.0	0.0	67
East South Central.....	5.9	0.0	362.3	47.5	30	17.8	112.9	0.0	71	0.0	0.0	36
West South Central.....	23.5	0.0	58.7	20.5	15	8.8	88.0	5.9	21	0.0	0.0	18
Mountain.....	0.0	0.0	442.2	24.1	1,544	16.1	130.7	8.0	370	0.0	8.0	193
Pacific.....	14.0	0.0	80.4	7.0	68	14.0	38.4	10.5	154	0.0	0.0	42
Total.....	10.7	0.2	119.9	11.4	268	20.6	87.8	3.8	162	0.0	2.3	68

TERRITORIES AND POSSESSIONS

Hawaii Territory

Honolulu—Dengue fever.—During the week ended December 11, 1943, 34 new cases of dengue fever were reported in Honolulu, T. H., bringing the total number of cases reported to date to 1,284.

FOREIGN REPORTS

CANADA

Provinces—Communicable diseases—Week ended November 27, 1943.—During the week ended November 27, 1943, cases of certain communicable diseases were reported by the Dominion Bureau of Statistics of Canada as follows:

Disease	Prince Edward Island	Nova Scotia	New Brunswick	Quebec	Ontario	Manitoba	Saskatchewan	Alberta	British Columbia	Total
Chickenpox		22		317	575	98	88	123	125	1,348
Diphtheria		13	3	64	1	4	3		7	95
Dysentery (amebic)					1				2	3
Dysentery (bacillary)				2						2
Encephalitis, infectious									1	1
German measles		1		3	16				9	29
Influenza	3	21	4		181	5			30	244
Measles		1		206	249	12	3	40	10	521
Meningitis, meningococcus			1		2	1				4
Mumps		7	2	76	196	36		27	83	432
Poliomyelitis					1	1				2
Scarlet fever		7	11	94	151	42	11	41	39	396
Tuberculosis (all forms)		6	2	70	59	6		29	41	213
Typhoid and paratyphoid fever				16	3				2	21
Undulant fever					5					5
Whooping cough		2		99	99	16	16	11	40	283

SWEDEN

Notifiable diseases—October 1943.—During the month of October 1943, cases of certain notifiable diseases were reported in Sweden as follows:

Disease	Cases	Disease	Cases
Cerebrospinal meningitis	8	Poliomyelitis	396
Diphtheria	177	Scarlet fever	3,064
Dysentery	162	Syphilis	114
Encephalitis, epidemic	4	Typhoid fever	4
Gonorrhea	2,023	Undulant fever	7
Hepatitis, epidemic	1,001	Well's disease	9
Paratyphoid fever	16		

(1936)

SWITZERLAND

Notifiable diseases—April-June 1943.—During the months of April, May, and June 1943, cases of certain notifiable diseases were reported in Switzerland as follows:

Disease	April	May	June
Cerebrospinal meningitis.....	11	4	7
Chickenpox.....	163	160	218
Diphtheria and croup.....	136	117	199
Dysentery.....	7	11	272
German measles.....	32	6	33
Hepatitis, epidemic.....	164	272	671
Influenza.....	8	8	35
Lethargic encephalitis.....			2
Malaria.....	1		
Measles.....	309	435	964
Mumps.....	172	158	158
Paratyphoid fever.....	3	9	20
Poliomyelitis.....	5	6	15
Scarlet fever.....	133	166	213
Trachoma.....	1	1	
Tuberculosis.....	402	404	483
Typhoid fever.....	2	6	11
Undulant fever.....	20	23	20
Whooping cough.....	141	253	673

WORLD DISTRIBUTION OF CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER

From medical officers of the Public Health Service, American consuls, International Office of Public Health, Pan American Sanitary Bureau, health section of the League of Nations, and other sources. The reports contained in the following tables must not be considered as complete or final as regards either the list of countries included or the figures for the particular countries for which reports are given.

CHOLERA

[C indicates cases]

NOTE.—Since many of the figures in the following tables are from weekly reports, the accumulated totals are for approximate dates.

Place	January-September 1943	October 1943	November 1943—week ended—			
			6	13	20	27
ASIA						
Ceylon.....	50					
China: Kwangsi Province.....	1,100					
India.....	210,136	32,897				
Bombay.....	16	12				
Calcutta.....	5,028	1,307	100	67	83	
Chittagong.....	245	37	36	34		
Cochin.....	192					
Madras.....	1,003	68	6	8		
Nagapatam.....	21					
Viragapatam.....	63					
Indis (French).....	55					
Chandernagor.....	8					
Karikal.....	30					
Pondichery.....	17					

¹ Cases reported up to Sept. 5, 1943, with a mortality rate of over 25 percent.

PLAGUE

[C indicates cases; D, deaths; P, present]

AFRICA						
Basutoland.....	C	23				
Belgian Congo.....	D		17			
Plague-infected rats.....	P					
British East Africa:						
Kenya.....	C	17				
Uganda.....	C	18				
Egypt ¹	C	14	1	3	12	
Port Said.....	C	6	1			
Madagascar.....	C	53				
Morocco (French).....	C	241	1	4		
Senegal.....	C	244				
Dakar.....	C	32				
Union of South Africa.....	C	65	1	2		

See footnotes at end of table.

PLAGUE—Continued

[C indicates cases; D, deaths; P, present]

Place		January- September 1943	October 1943	November 1943—week ended—			
				6	13	20	27
ASIA							
India.....	C	3,414	1,539	—	—	—	—
Indochina.....	C	30	1	—	—	—	—
Palestine.....	C	12	—	—	—	—	—
EUROPE							
Portugal (Azores). ⁴							
SOUTH AMERICA							
Ecuador: Loja Province.....	C	—	2	—	—	—	—
Peru:							
Lambayeque Department.....	C	2	—	—	—	—	—
Libertad Department.....	C	16	—	—	—	—	—
Lima Department.....	C	11	—	—	—	—	—
Lima.....	C	1	—	—	—	—	—
Plague-infected rats.....	P	—	—	—	—	—	—
Piura Department.....	C	2	—	—	—	—	—
Venezuela.....	C	10	—	—	—	—	—
OCEANIA							
Hawaii Territory:							
Hamakua District.....	D	5	—	—	—	—	—
Plague-infected rats.....		74	1	1	—	—	—

¹ Includes 12 pneumonic cases in a village south of Mafeteng.² Includes 7 suspected pneumonic plague deaths.³ A cablegram dated Dec. 13, 1943, states that 52 cases of plague have been reported to date in Egypt, including 47 cases at Suez, 4 at Port Tewfik, and 1 at Bitter Lake. For the week ended Dec. 13, 1 case was reported at Cairo, 2 at Port Said, 1 at El Bala, and 1 at Genefia.⁴ A report dated Nov. 19, 1943, states that during 1942 there were 54 cases of plague including 3 pneumonic cases and 2 septicemic cases among the civil population and 2 additional cases among the military population of the Azores. In 1943 the number of cases is about the same as for the year 1942.⁵ Includes 4 plague-infected mice.

SMALLPOX

[C indicates cases; D, deaths]

AFRICA							
Algeria.....	C	1,087	97				
Angola.....	C	613					
Basutoland.....	C	81					
Belgian Congo.....	C	2,975	435				
British East Africa:							
Kenya.....	C	1,488	448	120	145		
Mombasa.....	C	3					
Tanganyika.....	C	27	33	1			
Uganda.....	C		49	2	5		
Dahomey.....	C	141					
Egypt.....	C	2,848	456	67	44	61	
French Guinea.....	C	336	35				
Gold Coast.....	C	17	4				
Ivory Coast.....	C	144	10				
Mauritania.....	C	27	13				
Morocco (French).....	C	848					
Mozambique.....	C	1					
Nigeria.....	C	4,660	407	106	90	58	
Niger Territory.....	C	221	42				
Senegal.....	C	74					
Sierra Leone.....	C	3					
Sudan (French).....	C	3,440	156				
Tunisia.....	C	3					
Union of South Africa.....	C	439	2				
ASIA							
Arabia.....	C	1					
Ceylon.....	C	63	10	1	1		
India.....	C	34,428	3,275				
India (French).....	C	10					
Indochina.....	C	4,397	246				
Iran.....	C	510	11				
Iraq.....	C	195	33	6			
Palestine.....	C	101	8				
Syria and Lebanon.....	C	987	24	7	8		
Trans-Jordan.....	C	18	1				

SMALLPOX—Continued
[C indicates cases; D, deaths]

Place		January- September 1943	October 1943	November 1943—week ended—			
				6	13	20	27
EUROPE							
Belgium.....	C	1					
France.....	C	2					
Germany.....	C	1					
Gibraltar.....	C	1					
Portugal.....	C	40	2		2		
Scotland.....	C	1					1
Spain.....	C	204	5				
Switzerland.....	C	17					
Turkey.....	C	8, 161	499				
NORTH AMERICA							
British Honduras.....	C					1	
Canada.....	C	6					
Guatemala.....	C	27					
Mexico.....	C	306	21				
SOUTH AMERICA							
Brazil.....	C	44	5				
British Guiana.....	C	1					
Colombia.....	C	316	19	2	1	6	
Ecuador.....	C	18	4				
Peru.....	D	12					
Venezuela.....	C	91	4				

¹ On a vessel from North Africa.

TYPHUS FEVER

[C indicates cases; D, deaths]

AFRICA							
Algeria.....	C	8, 130	43				
Basutoland.....	C	18					
Belgian Congo.....	C	20	19				
British East Africa:							
Kenya.....	C	1	2	1			
Mombasa.....	C	1					
Uganda.....	C	1					
Egypt.....	C	39, 582	202	45	46	54	
Gold Coast.....	C	9					
Morocco (French).....	C	13, 552					
Morocco (Spanish).....	C	369					
Nigeria.....	C	9	2				
Portuguese East Africa.....	C		1				
Rhodesia, northern.....	C	10		4			
Senegal.....	C	2					
Dakar.....	C	15	4		3		
Sierra Leone.....	C	3					
Tunisia.....	C	232	20				
Union of South Africa.....	C	1, 587	8				
ASIA							
Afghanistan.....	C	520					
China: Shanghai.....	C	12					
India.....	C	1, 066					
Iran.....	C	9, 138					
Iraq.....	C	1, 421					
Palestine.....	C	266	30	9	1	6	8
Syria and Lebanon.....	C	81	8				
Trans-Jordan.....	C	15	2				
EUROPE							
Bulgaria.....	C	1, 712	33				
France—Seine Department.....	C	2					
Germany.....	C	1, 973					
Hungary.....	C	737	50		20	7	
Irish Free State.....	C	19			1		
Netherlands.....	C	1					
Portugal.....	C	9					
Rumania.....	C	6, 060	197				81
Slovakia.....	C	452	72	12	31		
Spain.....	C	578					
Turkey.....	C	3, 951					

¹ For the period Jan. 1 to Apr. 30, 1943.

TYPHUS FEVER—Continued

[C indicates cases; D, deaths]

Place	January- September 1943	October 1943	November 1943—week ended—			
			6	13	20	27
NORTH AMERICA						
Cuba.....	C	1				
Guatemala.....	C	967	145			
Jamaica.....	C	24	5		2	
Mexico.....	C	902	82			
SOUTH AMERICA						
Brazil.....	C	1				
Chile.....	C	211	9	2	2	1
Colombia.....	D	2				
Ecuador.....	C	277	42			
Peru.....	C	14				
Venezuela.....	C	17				
OCEANIA						
Australia.....	C	80	7	5	1	
Hawaii Territory.....	C	32	20	2	3	2

YELLOW FEVER

[C indicates cases; D, deaths]

AFRICA							
Belgian Congo:							
Bondo.....	D	2				1	
Kinshasa.....	D	1					
Leopoldville.....	C	2					
Stanleyville.....	D	1					
Yanonge.....	C	1					
British East Africa: Kenya—Misumu.....	C						11
Dahomey:							
Djougou District.....	C	12					
Natitingou.....	C	1					
French Guinea:							
Dubreka.....	D		1				
Matakang Island.....	D			1			
Gold Coast: Asubol.....	C	1					
Ivory Coast:							
Abidjan.....	C		1				
Toumodi.....	D				1		
Portuguese Guinea: ¹							
Senegal:							
Goudiri.....	D					1	
Kolda.....	C	1					
Tambacounda.....	C		1				
SOUTH AMERICA							
Brazil: Para State.....	D	1					
Colombia:							
Boyaca Department.....	D		4				
Cundinamarca Department.....	D	3					
Intendencia of Meta.....	D	2					
Santander Department.....	D		1				

¹ For the month of November 1943.² Suspected.³ Previously reported as having occurred at Conakry.⁴ During the week ended Nov. 13, 1943, a serious outbreak of yellow fever was reported in Portuguese Guinea. No figures are available.

X

FEDERAL SECURITY AGENCY
UNITED STATES PUBLIC HEALTH SERVICE

THOMAS PARRAN, *Surgeon General*

DIVISION OF PUBLIC HEALTH METHODS

G. St. J. PERROTT, *Chief of Division*

The PUBLIC HEALTH REPORTS, first published in 1878 under authority of an act of Congress of April 29 of that year, is issued weekly by the United States Public Health Service through the Division of Sanitary Reports and Statistics, pursuant to the following authority of law: United States Code, title 42, sections 7, 30, 93; title 44, section 220.

It contains (1) current information regarding the prevalence and geographic distribution of communicable diseases in the United States, insofar as data are obtainable, and of cholera, plague, smallpox, typhus fever, yellow fever, and other important communicable diseases throughout the world; (2) articles relating to the cause, prevention, and control of disease; (3) other pertinent information regarding sanitation and the conservation of the public health.

The PUBLIC HEALTH REPORTS is published primarily for distribution, in accordance with the law, to health officers, members of boards or departments of health, and other persons directly or indirectly engaged in public health work. Articles of special interest are issued as reprints or as supplements, in which forms they are made available for more economical and general distribution.


Requests for and communications regarding the PUBLIC HEALTH REPORTS, reprints, or supplements should be addressed to the Surgeon General, United States Public Health Service, Washington, D. C. Subscribers should remit direct to the Superintendent of Documents, Washington, D. C.

Librarians and others should preserve their copies for binding, as the Public Health Service is unable to supply the general demand for bound copies. Indexes will be supplied upon request.

UNITED STATES GOVERNMENT PRINTING OFFICE, WASHINGTON : 1943

For sale by the Superintendent of Documents, Washington, D. C.

Price 5 cents. Subscription price \$2.50 a year



PUBLIC LIBRARY
JAN 31 1944
DETROIT

Public Health Reports

Issued Weekly by the
UNITED STATES PUBLIC HEALTH SERVICE

Volume 58—Part 2

Numbers 27–53

July–December 1943



FEDERAL SECURITY AGENCY
UNITED STATES PUBLIC HEALTH SERVICE

THOMAS PARRAN, *Surgeon General*

DIVISION OF PUBLIC HEALTH METHODS

G. ST. J. PERROTT, *Chief of Division*

The PUBLIC HEALTH REPORTS, first published in 1878 under authority of an act of Congress of April 29 of that year, is issued weekly by the United States Public Health Service through the Division of Public Health Methods, pursuant to the following authority of law: United States Code, title 42, sections 7, 30, 93; title 44, section 220.

It contains (1) current information regarding the prevalence and geographic distribution of communicable diseases in the United States, insofar as data are obtainable, and of cholera, plague, smallpox, typhus fever, yellow fever, and other important communicable diseases throughout the world; (2) articles relating to the cause, prevention, and control of disease; (3) other pertinent information regarding sanitation and the conservation of the public health.

The PUBLIC HEALTH REPORTS is published primarily for distribution, in accordance with the law, to health officers, members of boards or departments of health, and other persons directly or indirectly engaged in public health work. Articles of special interest are issued as reprints or as supplements, in which forms they are made available for more economical and general distribution.

Requests for and communications regarding the PUBLIC HEALTH REPORTS, reprints, or supplements should be addressed to the Surgeon General, United States Public Health Service, Washington, D. C. Subscribers should remit direct to the Superintendent of Documents, Washington, D. C.

Librarians and others should preserve their copies for binding, as the Public Health Service is unable to supply the general demand for bound copies. Indexes will be supplied upon request.

UNITED STATES GOVERNMENT PRINTING OFFICE, WASHINGTON : 1944

For sale by the Superintendent of Documents, Washington 25, D. C.

Price 5 cents. Subscription price \$2.50 a year

INDEX

Key to Dates and Pages

No.	Date of issue	Pages	No.	Date of issue	Pages
27	July 2	1001-1032	41	Oct. 8	1497-1532
28	July 9	1033-1075	42	Oct. 15	1533-1572
29	July 16	1077-1119	43	Oct. 22	1573-1604
30	July 23	1121-1163	44	Oct. 29	1605-1640
31	July 30	1165-1199	45	Nov. 5	1641-1668
32	Aug. 6	1201-1231	46	Nov. 12	1669-1700
33	Aug. 13	1233-1264	47	Nov. 19	1701-1728
34	Aug. 20	1265-1292	48	Nov. 26	1729-1760
35	Aug. 27	1293-1327	49	Dec. 3	1761-1792
36	Sept. 3	1328-1364	50	Dec. 10	1793-1836
37	Sept. 10	1365-1392	51	Dec. 17	1837-1880
38	Sept. 17	1393-1428	52	Dec. 24	1881-1907
39	Sept. 24	1429-1456	53	Dec. 31	1909-1940
40	Oct. 1	1457-1496			

A

	Page
Absenteeism, sickness, among industrial workers, first quarter of 1943, with an inquiry into the occurrence of the respiratory diseases, 1934-43 [Gafafer].....	1273
Absenteeism, sickness, among industrial workers, second quarter of 1943, with a note on the occurrence of the respiratory diseases, 1934-43 [Gafafer].....	1885
Absenteeism, sickness, among male and female industrial workers, 1933-42, inclusive [Gafafer].....	1250
Actinomycosis. (See United States—Monthly State reports.)	
<i>Aerobacter cloacae</i> , studies on strains of, responsible for acute illness among workers using low-grade stained cotton [Caminita, Schreiner, Kolb, and Neal].....	1165
Alabama. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Algeria: Infectious diseases—May–August 1943.....	1791
Altman, Isidore: The patient load of physicians in private practice. A comparative statistical study of three areas.....	1329
<i>Amblyomma americanum</i> , the occurrence of <i>Rickettsia diaporica</i> in, in eastern Texas. American Q fever: [Parker and Kohls].....	1510
Angola: Notifiable diseases.....	1423, 1877
<i>Anopheles</i> , Puerto Rican, identification of first stage larvae of [Pratt].....	1715
Anthrax: (See also United States—Monthly State reports.)	
Weekly city reports.....	1117, 1226,
1360, 1389, 1452, 1568, 1599, 1662, 1697, 1725, 1755, 1876	
Weekly State reports.....	1024,
1068, 1114, 1156, 1190, 1224, 1258, 1286, 1319, 1358, 1386, 1418,	
1449, 1489, 1520, 1565, 1596, 1634, 1660, 1694, 1722, 1752, 1783,	
1830, 1873, 1901, 1932	
Arizona. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Arkansas. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Armstrong, Wallace D.: The effect of topically applied sodium fluoride on dental caries experience.....	1701
Arsenic in water contaminated with chemical warfare agents, the detection and analysis of [Ruchhoff, Placak, and Schott].....	1761
Atabrine and sulfadiazine, toxic effects of, in growing rats [Wright and Lillie].....	1242

B

<i>Bacillus pestis</i> cultures, twenty-year survival of virulent, without transfer [Francis].....	1379
--	------

	Page
<i>Balantidium coli</i> infections, carbarsone treatment for [Young and Burrows].	1272
Baum, William S.: The diet in Germany and the occupied countries during the second world war.	1669
Black, Luther A.: Surveys of milk laboratories in war areas in the United States—	
I. Practices observed in making agar plate counts.	1605
II. Practices observed in making direct microscopic examinations and methylene blue reduction tests.	1641
III. Observations on sampling and health department practice relative to bacteriological milk analysis.	1681
Blood calcium, effect of lead absorption on [Jenrette and Fairhall].	1001
Botulism. (See United States—Monthly State reports.)	
Botvinick, Isadore: A soap which indicates the presence of mercury fulminate.	1183
Brazil:	
Para State—Belém—Poliomyelitis.	1363
Rio Grande do Sul State—Poliomyelitis.	1530
Brazil, Colombia, and the United States:	
Experimental transmission of the rickettsiae of the spotted fevers of, by the argasid tick <i>Ornithodoros nicolleti</i> [Davis].	1742
Experimental transmission of the spotted fevers of, by the argasid tick <i>Ornithodoros parkeri</i> [Davis].	1201
The tick <i>Ornithodoros rudis</i> as a host to the rickettsiae of the spotted fevers of [Davis].	1016
British East Africa: Tanganyika Territory—Cerebrospinal meningitis.	1791
Burns and shock, experimental chemotherapy of. IV. Production of traumatic shock in mice. V. Therapy with mouse serum and sodium salts [Rosenthal].	1429
Burrows, Robert: Carbarsone treatment for <i>Balantidium coli</i> infections.	1272
Butterfield, C. T.: Influence of pH and temperature on the survival of coliforms and enteric pathogens when exposed to free chlorine.	1837
C.	
California: (See also Name of disease—Weekly State reports; United States—Monthly State reports.)	
Plague infection in.	1361,
1389, 1422, 1493, 1528, 1529, 1569, 1599, 1600, 1635, 1697,	1876
Caminita, B. H.: Studies on strains of <i>Aerobacter cloacae</i> responsible for acute illness among workers using low-grade stained cotton.	1165
Canada: Provinces—Communicable diseases.	1029,
1073, 1118, 1161, 1194, 1228, 1263, 1291, 1324, 1363, 1390, 1423, 1453,	
1494, 1530, 1570, 1600, 1637, 1664, 1698, 1727, 1756, 1791, 1834, 1877,	
1906, 1936	
Cancer of the lung, the incidence and prevalence of [Dorn].	1265
Carbarsone treatment for <i>Balantidium coli</i> infections [Young and Burrows].	1272
Chambers, C. W.: Influence of pH and temperature on the survival of coliforms and enteric pathogens when exposed to free chlorine.	1837
Chickenpox. (See United States—Monthly State reports.)	
Chlorine, free, influence of pH and temperature on the survival of coliforms and enteric pathogens when exposed to [Butterfield, Wattie, Megregian, and Chambers].	1837
Cholera: Foreign reports.	1162, 1195, 1229, 1325, 1454, 1495, 1638, 1757, 1937
Ciocco, Antonio: The patient load of physicians in private practice. A comparative statistical study of three areas.	1329
<i>Clostridium perfringens</i> (Welchii) infections in guinea pigs, the mechanism of antitoxic immunity in [Stewart].	1277
Coccidioidomycosis. (See United States—Monthly State reports.)	
Coliforms and enteric pathogens, influence of pH and temperature on the survival of, when exposed to free chlorine [Butterfield, Wattie, Megregian, and Chambers].	1837
Collins, Selwyn D.: Extent of immunization and case histories for diphtheria, smallpox, scarlet fever, and typhoid fever in 200,000 surveyed families in 28 large cities.	1121
Colombia, Brazil, and the United States:	
Experimental transmission of the rickettsiae of the spotted fevers of, by the argasid tick <i>Ornithodoros nicolleti</i> [Davis].	1742

Colombia, Brazil, and the United States—Continued.	Page
Experimental transmission of the spotted fevers of, by the argasid tick <i>Ornithodoros parkeri</i> [Davis].....	1201
The tick <i>Ornithodoros rudis</i> as a host to the rickettsiae of the spotted fevers of [Davis].....	1016
Colorado: (See also Name of disease—Weekly State reports; United States—Monthly State reports.)	
Plague infection in.....	1160, 1290, 1422
Communicable diseases. (See Name of disease; Name of country.)	
Community services vs. lost man-hours [McNutt].....	1082
Conference, forty-first annual, of the United States Public Health Service with the State and Territorial health officers:	
Abstracts of committee reports adopted by the.....	1101
Opening remarks to the [Parran].....	1077
Conjunctivitis. (See United States—Monthly State reports.)	
Connecticut. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Cooper, Marcia: A survey of statistical studies on the prevalence and incidence of mental disorder in sample populations.....	1909
Councell, Clara: Extent of immunization and case histories for diphtheria, smallpox, scarlet fever, and typhoid fever in 200,000 surveyed families in 28 large cities.....	1121
Court decisions on public health:	
Children bitten by rats—action against owners of adjoining premises ..	1427
City ordinance regarding maintenance of clean and habitable premises upheld.....	1532
City tourist camp ordinance held to conflict with State trailer park law.....	1867
Death certificates—statements as to cause of death.....	1230
Death from lead poisoning held compensable under workmen's compensation act.....	1031
Garbage—dumping—construction of statute concerning.....	1718
Garbage—ordinance making collection by unauthorized person unlawful upheld.....	1571
Inflammation of the eyes of the newborn—reporting.....	1031
Liability of physician for failure to use prophylactic in infant's eyes at birth.....	1199
Piggery—held nuisance—operation enjoined.....	1496
Public health law construed.....	1868
Safe drinking water at State institution—liability of State officials for failure to furnish.....	1230
Public water supply—protection against pollution—reasonableness of State board of health regulations.....	1666
Sewage disposal—stream pollution by city—order of State stream control commission upheld.....	1198
State sanitary code amendment defining milk products held invalid as applied to certain product.....	1700
Swine—keeping for feeding on swill, etc., brought from without town where animals are kept.....	1779
Typhoid fever—held to be accidental injury under workmen's compensation law.....	1030
Typhus fever—contraction by bakery employee—liability of employer.....	1426
Uniform narcotic drug act—earlier law repealed by—marihuana included under.....	1868
Unwholesome food—sale.....	1152
Venereal disease—exposing another person to.....	1604
Cuba:	
Habana—Communicable diseases.....	1194, 1228, 1570, 1601, 1637, 1664, 1878
Provinces—Notifiable diseases.....	1161, 1228, 1324, 1494, 1601, 1664, 1878
Sagua La Grande—Typhoid fever.....	1424

D

Daft, Floyd S.: The successful treatment of granulocytopenia and leukopenia in rats with crystalline folic acid.....	1542
Davis, Burnet M.: War and the distribution of physicians.....	1545
Davis, Dorland J.: Infection in monkeys with strains of <i>Trypanosoma cruzi</i> isolated in the United States.....	1006

	Page
Davis, Gordon E.: Experimental transmission of the rickettsiae of the spotted fevers of Brazil, Colombia, and the United States by the argasid tick <i>Orni-</i> <i>thodoros nicolleti</i>	1742
Experimental transmission of the spotted fevers of the United States, Colombia, and Brazil by the argasid tick <i>Ornithodoros parkeri</i>	1201
The tick <i>Ornithodoros rudis</i> as a host to the rickettsiae of the spotted fevers of Colombia, Brazil, and the United States.....	1016
Deaths: (See also Mortality.)	
Large cities—Weekly reports (Weekly Mortality Index)--	1020, 1064, 1110,
1152, 1186, 1220, 1254, 1292, 1314, 1354, 1382, 1426, 1439, 1485, 1516,	
1561, 1592, 1636, 1666, 1690, 1718, 1748, 1778, 1825, 1869, 1896, 1927	
Delaware. (See Name of disease—Weekly State reports; United States— Monthly State reports.)	
Dengue fever: (See also United States—Monthly State reports.)	
Hawaii Territory—Honolulu.....	1290, 1322,
1422, 1529, 1569, 1600, 1663, 1697, 1726, 1833, 1904, 1935	
In Honolulu.....	1823
Dental caries experience, effect of topically applied sodium fluoride on [Knutson and Armstrong].....	1701
<i>Demacantor andersoni</i> Stiles, <i>Salmonella enteritidis</i> : experimental trans- mission by the Rocky Mountain wood tick [Parker and Steinhause].....	1010
Dermatitis, an outbreak of, from hair lacquer [Schwartz].....	1623
Diarrhea and enteritis. (See United States—Monthly State reports.)	
Diarrhea, infant, outbreak of, in a Maine hospital.....	1560
Diet in Germany and the occupied countries during the second world war [Spicknall, Fishburn, and Baum].....	1669
Dinan, J. F.: The promin treatment of leprosy. A progress report.....	1729
Diphtheria: (See also United States—Monthly State reports.)	
Weekly city reports.....	1025, 1069, 1115, 1157, 1191, 1225,
1259, 1287, 1320, 1359, 1387, 1419, 1450, 1490, 1526, 1566, 1597,	
1633, 1661, 1695, 1723, 1753, 1788, 1831, 1874, 1902, 1933	
Weekly State reports.....	1022, 1066, 1112, 1154, 1188, 1222,
1256, 1284, 1317, 1356, 1384, 1416, 1447, 1487, 1518, 1563, 1594,	
1630, 1658, 1692, 1720, 1750, 1781, 1823, 1871, 1899, 1930	
Diphtheria, smallpox, scarlet fever, and typhoid fever, extent of im- munization and case histories for, in 200,000 surveyed families in 28 large cities [Collins and Council].....	1121
Disabilities causing absence from work, frequency and duration of, among the employees of a public utility, 1938-42 [Gafafer].....	1554
District of Columbia. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Dog bites. (See United States—Monthly State reports.)	
Dominican Republic:	
Influenza.....	1792
Malaria.....	1792
Dorn, Harold F.: The incidence and prevalence of cancer of the lung.....	1265
Dysentery: (See also United States—Monthly State reports.)	
Weekly city reports.....	1027, 1071, 1117, 1159, 1192,
1226, 1261, 1289, 1322, 1360, 1389, 1421, 1452, 1492, 1528, 1568,	
1599, 1634, 1662, 1697, 1725, 1755, 1790, 1833, 1876, 1904, 1935	
Weekly State reports.....	1024, 1068, 1114, 1156, 1190, 1224,
1258, 1286, 1319, 1358, 1386, 1418, 1449, 1489, 1520, 1565, 1596,	
1632, 1660, 1694, 1722, 1752, 1783, 1830, 1873, 1901, 1932	

E

Eccles, C. G.: The promin treatment of leprosy. A progress report.....	1729
Egypt:	
Infectious diseases—First quarter 1943.....	1834
Vital statistics—First quarter 1943.....	1834
Encephalitis, infectious: (See also United States—Monthly State reports.)	
Weekly city reports.....	1025, 1069, 1115, 1157, 1191,
1225, 1259, 1287, 1320, 1359, 1387, 1419, 1450, 1490, 1526, 1566,	
1597, 1633, 1661, 1695, 1723, 1753, 1788, 1831, 1874, 1902, 1933	
Weekly State reports.....	1024, 1068, 1114, 1156, 1190,
1224, 1258, 1286, 1319, 1358, 1386, 1418, 1449, 1489, 1520, 1565,	
1596, 1632, 1660, 1694, 1722, 1752, 1783, 1830, 1873, 1901, 1932	

F

	Page
Faget, G. H.: The promin treatment of leprosy. A progress report.....	1729
Fairhall, Lawrence T.: Effect of lead absorption on blood calcium.....	1001
Finland—Notifiable diseases.....	1570, 1665, 1878
Fishburn, Howard D.: The diet in Germany and the occupied countries during the second world war.....	1669
Florida. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Folic acid, crystalline, the successful treatment of granulocytopenia and leukopenia in rats with [Daft and Sebrell].....	1542
Food poisoning. (See United States—Monthly State reports.)	
Food poisoning, study of an outbreak of, in a hospital in Galveston, Texas [Lumsden, Nau, and Stead].....	1497
Foreign reports: Quarantinable diseases.....	1029, 1074, 1118, 1162, 1195, 1229, 1264, 1291, 1325, 1363, 1391, 1425, 1454, 1495, 1531, 1571, 1603, 1638, 1665, 1698, 1728, 1757, 1792, 1836, 1879, 1906, 1927
Francis, Edward: Twenty-year survival of virulent <i>Bacillus pestis</i> cultures without transfer.....	1379

G

Gafafer, W. M.:	
Frequency and duration of disabilities causing absence from work among the employees of a public utility.....	1554
Sickness absenteeism among industrial workers, first quarter of 1943, with an inquiry into the occurrence of the respiratory diseases, 1934-43.....	1273
Sickness absenteeism among industrial workers, second quarter of 1943, with a note on the occurrence of the respiratory diseases, 1934-43..	1885
Sickness absenteeism among male and female industrial workers, 1933-42, inclusive.....	1250
Georgia. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
German measles. (See United States—Monthly State reports.)	
Germany: Infectious diseases—	
Week ended July 31, 1943, and January 1 to July 25, 1943—Comparative.....	1424
Week ended September 25, 1943, and January 1 to September 18, 1943—Comparative.....	1835
Germany and the occupied countries, the diet in, during the second world war [Spicknall, Fishburn, and Baum].....	1669
Gilliam, Alexander G.: Jaundice following administration of human serum.....	1233
Gover, Mary: Influenza and pneumonia mortality in a group of 90 cities in the United States, August 1935-March 1943, with a summary for August 1920-March 1943.....	1033
Granulocytopenia and leukopenia, the successful treatment of, in rats with crystalline folic acid [Daft and Sebrell].....	1542
Granuloma. (See United States—Monthly State reports.)	
Great Britain—England and Wales: Infectious diseases—	
Years 1940, 1941, and 1942—Comparative.....	1602
13 weeks ended April 3, 1943, and July 3, 1943.....	1756

H

Hair lacquer, an outbreak of dermatitis from [Schwartz].....	1623
Hampton, Brock C.: Smallpox in relation to State vaccination laws and regulations.....	1771
Haugen, Gerhard B.: An approach to the mental hygiene-public health problem.....	1211
Hawaii Territory:	
Honolulu—	
Dengue fever.....	1290,
1322, 1422, 1529, 1569, 1600, 1663, 1697, 1726, 1833, 1904, 1935	
Outbreak of mild influenza.....	1117
Plague (human).....	1422
Plague (rodent).....	1028, 1072, 1262, 1529, 1726, 1755, 1904
Health education in 1943, fellowships in.....	1214
Health of the United States at war.....	1281

	Page
Heilig, R.: A note on the rickettsioses in India.....	1208
Hilleboe, Herman E.: Opportunities in the newer methods of tuberculosis case finding.....	1094
Tuberculosis mortality in the United States, 1939-41.....	1457
Honolulu, dengue fever in.....	1823
Hookworm disease. (See United States—Monthly State reports.)	
Hospitalization, incidence of.....	1220, 1314, 1690, 1778

I

Idaho. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Illinois. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Immunization, extent of, and case histories for diphtheria, smallpox, scarlet fever, and typhoid fever in 200,000 surveyed families in 28 large cities [Collins and Council]	1121
Impetigo contagiosa. (See United States—Monthly State reports.)	
India, a note on the rickettsioses in [Topping, Heilig, and Naidu].....	1208
Indiana. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Infantile paralysis. (See Poliomyelitis.)	
Infectious diseases. (See Name of country.)	
Influenza: (See also United States—Monthly State reports.)	
Weekly city reports.....	1025, 1069, 1115, 1157, 1191, 1225, 1259, 1287, 1320, 1359, 1387, 1419, 1450, 1490, 1526, 1566, 1597, 1633, 1661, 1695, 1723, 1753, 1788, 1831, 1874, 1902, 1933
Weekly State reports.....	1022, 1066, 1112, 1154, 1188, 1222, 1256, 1284, 1317, 1356, 1384, 1416, 1447, 1487, 1518, 1563, 1594, 1630, 1658, 1692, 1720, 1750, 1781, 1828, 1871, 1899, 1930
Influenza and pneumonia mortality in a group of 90 cities in the United States, August 1935-March 1943, with a summary for August 1920-March 1943 [Gover].....	1033
Influenza, outbreak of mild, in Honolulu, Hawaii.....	1117
Iowa. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Irish Free State: Vital statistics—First quarter ended March 31, 1943....	1162

J

Jacobs, Leon: Studies on trichinosis. XV. Summary of the findings of <i>Trichinella spiralis</i> in a random sampling and other samplings of the population of the United States.....	1293
Jamaica: Communicable diseases.....	1073, 1194, 1325, 1453, 1603, 1727, 1879
Jaundice. (See United States—Monthly State reports.)	
Jaundice following administration of human serum [Oliphant, Gilliam, and Larson].....	1233
Jenrette, Wendell V.: Effect of lead absorption on blood calcium.....	1001
Johansen, F. A.: The promin treatment of leprosy. A progress report...	1729

K

	Page
Kansas. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Karpinos, Bernard D.: The physically handicapped.....	1573
Kentucky. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Kerr, K. B.: Studies on trichinosis. XV. Summary of the findings of <i>Trichinella spiralis</i> in a random sampling and other samplings of the population of the United States.....	1293
Knutson, John W.: The effect of topically applied sodium fluoride on dental caries experience.....	1701
Kohls, Glen M.: American Q fever: the occurrence of <i>Rickettsia diaporica</i> in <i>Amblyomma americanum</i> in eastern Texas.....	1510
Kolb, R. W.: Studies on strains of <i>Aerobacter cloacae</i> responsible for acute illness among workers using low-grade stained cotton.....	1165

L

	Page
Larson, Carl L.: Jaundice following administration of human serum.....	1233
Lead absorption, effect of, on blood calcium [Jenrette and Fairhall].....	1001
Lemkau, Paul: A survey of statistical studies on the prevalence and incidence of mental disorder in sample populations.....	1909
Leprosy: (See also United States—Monthly State reports.)	
Weekly city reports.....	1159, 1261, 1599, 1833, 1935
Weekly State reports.....	1024,
1068, 1114, 1156, 1190, 1224, 1258, 1286, 1319, 1358, 1386, 1418,	
1449, 1489, 1520, 1565, 1596, 1632, 1660, 1694, 1722, 1752, 1783,	
1830, 1873, 1901, 1932	
Leprosy, the promin treatment of. A progress report. [Faget, Pogge, Johansen, Dinan, Prejean, and Eccles].....	1729
Leukopenia and granulocytopenia, the successful treatment of, in rats with crystalline folic acid [Daft and Sebrell].....	1542
Lillie, R. D.: Notes on the pathology of experimental trinitrotoluene poisoning....	1436
Toxic effects of atabrine and sulfadiazine in growing rats.....	1242
Louisiana. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Lumsden, L. L.: Study of an outbreak of food poisoning in a hospital in Galveston, Texas.....	1497
Lymphocytic choriomeningitis. (See United States—Monthly State reports.)	
Lymphogranuloma venereum. (See United States—Monthly State reports.)	

Mc

McNutt, Paul V.: Community services vs. lost man-hours.....	1082
---	------

M

Maine. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Malaria. (See United States—Monthly State reports.)	
Martinique: Fort de France—Influenza.....	1757
Maryland. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Mason, Howard S.: A soap which indicates the presence of mercury fulminate.....	1183
Massachusetts. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Measles: (See also United States—Monthly State reports.)	
Weekly city reports.....	1025, 1069, 1115, 1157, 1191,
1225, 1259, 1287, 1320, 1359, 1387, 1419, 1450, 1490, 1526, 1566,	
1597, 1633, 1661, 1695, 1723, 1753, 1788, 1831, 1874, 1902, 1933	
Weekly State reports.....	1022, 1066, 1112, 1154, 1188,
1222, 1256, 1284, 1317, 1356, 1384, 1416, 1447, 1487, 1518, 1563,	
1594, 1630, 1658, 1692, 1720, 1750, 1781, 1828, 1871, 1899, 1930	
Megregian, Stephen: Influence of pH and temperature on the survival of coliforms and enteric pathogens when exposed to free chlorine.....	1837
Meningitis, meningococcus: (See also United States—Monthly State reports.)	
Weekly city reports.....	1025, 1069, 1115, 1157, 1191,
1225, 1259, 1287, 1320, 1359, 1387, 1419, 1450, 1490, 1526, 1566,	
1597, 1633, 1661, 1695, 1723, 1753, 1788, 1831, 1874, 1902, 1933	
Weekly State reports.....	1022, 1066, 1112, 1154, 1188,
1222, 1256, 1284, 1317, 1356, 1384, 1416, 1447, 1487, 1518, 1563,	
1594, 1630, 1658, 1692, 1720, 1750, 1781, 1828, 1871, 1899, 1930	
Mental disorder in sample populations, a survey of statistical studies on the prevalence and incidence of [Lemkau, Tietze, and Cooper].....	1909
Mental hygiene-public health problem, an approach to the [Haugen].....	1211
Mercury fulminate, a soap which indicates the presence of [Mason and Botvinick].....	1183
Mexico:	
Cananea—Influenza.....	1835
Vera Cruz—Dengue fever.....	1835

	Page
Michigan. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Milk laboratories, surveys of, in war areas in the United States [Black]:	
I. Practices observed in making agar plate counts.....	1605
II. Practices observed in making direct microscopic examinations and methylene blue reduction tests.....	1641
III. Observations on sampling and health department practices relative to bacteriological milk analysis.....	1681
Milmore, B. K.: Harborage of <i>Rattus rattus alexandrinus</i>	1507
Minnesota. (See Name of disease—Weekly State reports, United States—Monthly State reports.)	
Mississippi. (See Name of disease—Weekly State reports, United States—Monthly State reports.)	
Missouri. (See Name of disease—Weekly State reports, United States—Monthly State reports.)	
Montana: (See also Name of disease—Weekly State reports, United States—Monthly State reports.)	
Plague infection in.....	1362, 1422, 1493, 1569, 1600
Morgan, Russell H.: The automatic control of exposure in photofluorography.....	1533
Morocco: Infectious diseases—May–August 1943.....	1792
Mortality: (See also Deaths.)	
Influenza and pneumonia, in a group of 90 cities in the United States, August 1935–March 1943, with a summary for August 1920–March 1943 [Gover].....	1033
Tuberculosis, in the United States, 1939–41 [Yerushalmy, Hilleboe, and Palmer].....	1457
Mortality rates, provisional, for the first quarter of 1943.....	1439
Mountin, J. W.: Outlook for the coming year.....	1088
Mumps. (See United States—Monthly State reports.)	
Munitions manufacturing, surveys of liquid wastes from:	
I. Trinitrotoluene (TNT) wastes [Smith and Walker].....	1365
II. Smokeless powder wastes [Smith and Walker].....	1373
III. Small arms ammunition [Smith and Walker].....	1393
IV. Tetryl wastes [Smith].....	1404
V. Nitroglycerine wastes [Smith].....	1409
N	
Naidu, V. R.: A note on the rickettsioses in India.....	1208
Nau, C. A.: Study of an outbreak of food poisoning in a hospital in Galveston, Texas.....	1497
Neal, P. A.: Studies on strains of <i>Aerobacter cloacae</i> responsible for acute illness among workers using low-grade stained cotton.....	1165
Nebraska. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Nevada. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
New Hampshire. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
New Jersey. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
New Mexico: (See also Name of disease—Weekly State reports; United States—Monthly State reports.)	
Plague infection in.....	1027, 1071
New York. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
New Zealand:	
Hospital ship, smallpox on a.....	1073
Vital statistics—Year 1942–43.....	1390
Nicaragua: Poliomyelitis.....	1835
North Carolina. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
North Dakota. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Norway: Trondheim—Diphtheria.....	1727
Notifiable diseases. (See Name of country.)	

O

Page

Ohio. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Oklahoma. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Oliphant, John W.: Jaundice following administration of human serum..	1233
Ophthalmia neonatorum. (See United States—Monthly State reports.)	
<i>Opisocrostis bruneri</i> (Baker) and <i>Thrassis bacchi</i> (Roths.), report on the fleas, as vectors of plague [Prince].....	1013
Oregon: (See also Name of disease—Weekly State reports; United States—Monthly State reports.)	
Plague infection in	1028, 1160, 1261
<i>Ornithodoros nicolleti</i> , the argasid tick, experimental transmission of the rickettsiae of the spotted fevers of Brazil, Colombia, and the United States by [Davis].....	1742
<i>Ornithodoros parkeri</i> , the argasid tick, experimental transmission of the spotted fevers of the United States, Colombia, and Brazil by [Davis].....	1201
<i>Ornithodoros rudis</i> , the tick, as a host to the rickettsiae of the spotted fevers of Colombia, Brazil, and the United States [Davis].....	1016
Outlook for the coming year [Mountin].....	1088

P

Palmer, C. E.: Tuberculosis mortality in the United States, 1939-41	1457
Panama Canal Zone: Notifiable diseases..	1072, 1227, 1323, 1600, 1636, 1726, 1905
Paraguay: Asuncion—Poliomyelitis.....	1906
Paratyphoid fever. (See typhoid fever.)	
Parker, R. R.: American Q fever: the occurrence of <i>Rickettsia diaporica</i> in <i>Amblyomma americanum</i> in eastern Texas.....	1510
<i>Salmonella enteritidis</i> : experimental transmission by the Rocky Mountain wood tick <i>Dermacentor andersoni</i> Stiles.....	1010
Parran, Thomas: Opening remarks to the forty-first annual conference of the United States Public Health Service with the State and Territorial health officers.....	1077
Patient load of physicians in private practice. A comparative statistical study of three areas [Ciocco and Altman].....	1329
Pellagra. (See United States—Monthly State reports.)	
Pennsylvania. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Perrott, G. St. J.: War and the distribution of physicians.....	1545
Photofluorography, the automatic control of exposure in [Morgan].....	1533
Physically handicapped, the [Karpinos].....	1573
Physicians, the patient load of, in private practice. A comparative statistical study of three areas [Ciocco and Altman].....	1329
Physicians, war and the distribution of [Perrott and Davis].....	1545
Placak, O. R.: The detection and analysis of arsenic in water contaminated with chemical warfare agents.....	1761
Plague: (See also United States—Monthly State reports.)	
California—Siskiyou County (human case).....	1361
Foreign reports.....	1074,
1118, 1162, 1195, 1229, 1264, 1291, 1325, 1363, 1391, 1425, 1454, 1495, 1531, 1571, 1603, 1638, 1698, 1728, 1758, 1836, 1879, 1906, 1937	
Hawaii Territory—	
Human.....	1422
Rodent.....	1028, 1072, 1262, 1529, 1726, 1755, 1904
Plague, vectors of, report on the fleas <i>Opisocrostis bruneri</i> (Baker) and <i>Thrassis bacchi</i> (Roths.) as [Prince].....	1013
Plague infection:	
California.....	1361,
1389, 1422, 1493, 1528, 1529, 1569, 1599, 1600, 1635, 1697, 1876	
Colorado.....	1160, 1290, 1422
Montana.....	1362, 1422, 1493, 1569, 1600
New Mexico.....	1027, 1071
Oregon.....	1028, 1160, 1261
Wyoming.....	1290, 1600

	Page
Pneumonia (all forms): (See also United States—Monthly State reports.)	
Weekly city reports.....	1025,
1069, 1115, 1157, 1191, 1225, 1259, 1287, 1320, 1359, 1387, 1419,	
1450, 1490, 1526, 1566, 1597, 1633, 1661, 1695, 1723, 1753,	
1788, 1831, 1874, 1902, 1933	
Pneumonia and influenza mortality in a group of 90 cities in the United States, August 1935–March 1943, with a summary for August 1920–March 1943 [Gover].....	1033
Pogge, R. C.: The promin treatment of leprosy. A progress report.....	1729
Poliomylitis: (See also United States—Monthly State reports.)	
Brazil—Para State—Belém.....	1363
Prevalence of.....	1108, 1412
Weekly city reports.....	1025,
1069, 1115, 1157, 1191, 1225, 1259, 1287, 1320, 1359, 1387, 1419,	
1450, 1490, 1526, 1566, 1597, 1633, 1661, 1695, 1723, 1753, 1788,	
1831, 1874, 1902, 1933	
Weekly State reports.....	1023,
1067, 1113, 1155, 1189, 1223, 1257, 1285, 1318, 1357, 1385, 1417,	
1448, 1488, 1519, 1564, 1595, 1631, 1659, 1693, 1721, 1751,	
1782, 1829, 1872, 1900, 1931	
Porges, Ralph: The use of curtain walls in ratproofing.....	1881
Pratt, Harry D.: The identification of first stage larvae of Puerto Rican <i>Anopheles</i>	1715
Prejean, B. M.: The promin treatment of leprosy. A progress report..	1729
Prince, Frank M.: Report on the fleas <i>Opisocrostis bruneri</i> (Baker) and <i>Thrassius bacchi</i> (Roths.) as vectors of plague.....	1013
Promin treatment of leprosy. A progress report. [Faget, Pogge, Johansen, Dinan, Prejean, and Eccles].....	1729
Psittacosis. (See United States—Monthly State reports.)	
Public health, court decisions on. (See Court decisions on public health.)	
Public health-mental hygiene problem, an approach to the [Haugen]....	1211
Public Health Service publications. A list of publications issued during the period January–June 1943.....	1511
Puerperal septicemia. (See United States—Monthly States reports.)	
Puerto Rico: Influenza.....	1493

Q

Q fever, American: the occurrence of <i>Rickettsia diaporica</i> in <i>Amblyomma americanum</i> in eastern Texas [Parker and Kohls].....	1510
--	------

R

Rabies:	
In animals. (See United States—Monthly State reports.)	
In man. (See United States—Monthly State reports.)	
Rat-bite fever. (See United States—Monthly State reports.)	
Ratproof construction of vessels, the benefits accruing from the [Sherrard]..	1888
Ratproofing, the use of curtain walls in [Porges].....	1881
<i>Rattus rattus alexandrinus</i> , harborage of [Milmore].....	1507
Relapsing fever. (See United States—Monthly State reports.)	
Respiratory diseases, 1934–43, sickness absenteeism among industrial workers, first quarter of 1943, with a note on the occurrence of the [Gafafer].....	1273
Respiratory diseases, 1934–43, sickness absenteeism among industrial workers, second quarter of 1943, with a note on the occurrence of the [Gafafer].....	1885
Rheumatic fever. (See United States—Monthly State reports.)	
Rhode Island. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
<i>Rickettsia diaporica</i> , the occurrence of, in <i>Amblyomma americanum</i> in eastern Texas. American Q fever: [Parker and Kohls].....	1510
Rickettsioses in India, a note on the [Topping, Heilig, and Naidu].....	1208

	Page
Rocky Mountain spotted fever: (<i>See also</i> United States—Monthly State reports.)	
Weekly city reports.....	1027, 1071, 1117, 1159, 1261, 1322, 1360, 1389, 1421, 1452, 1528, 1599, 1662
Weekly State reports.....	1024, 1068, 1114, 1156, 1190, 1224, 1258, 1286, 1319, 1258, 1386, 1418, 1449, 1489, 1520, 1565, 1596, 1632, 1660, 1694, 1722, 1752, 1783, 1830, 1873, 1901, 1932
Rosenthal, Sanford M.: Experimental chemotherapy of burns and shock. IV. Production of traumatic shock in mice. V. Therapy with mouse serum and sodium salts.....	1429
Ruchhoft, C. C.: The detection and analysis of arsenic in water contaminated with chemical warfare agents.....	1761

S

<i>Salmonella enteritidis</i> : experimental transmission by the Rocky Mountain wood tick <i>Dermacentor andersoni</i> Stiles [Parker and Steinhaus].....	1010
Sanitation standards, emergency minimum.....	1793
Scarlet fever: (<i>See also</i> United States—Monthly State reports.)	
Weekly city reports.....	1025, 1069, 1115, 1157, 1191, 1225, 1259, 1287, 1320, 1359, 1387, 1419, 1450, 1490, 1526, 1566, 1597, 1633, 1661, 1695, 1723, 1753, 1788, 1831, 1874, 1902, 1933
Weekly State reports.....	1023, 1067, 1113, 1155, 1189, 1223, 1257, 1285, 1318, 1357, 1385, 1417, 1448, 1488, 1519, 1564, 1595, 1631, 1659, 1693, 1721, 1751, 1782, 1829, 1872, 1900, 1931
Scarlet fever, diphtheria, smallpox, and typhoid fever, extent of immunization and case histories for, in 200,000 surveyed families in 28 large cities [Collins and Council].....	1121
Schneider, R.: Studies on strains of <i>Aerobacter cloacae</i> responsible for acute illness among workers using low-grade stained cotton.....	1165
Schott, Stuart: The detection and analysis of arsenic in water contaminated with chemical warfare agents.....	1761
Schwartz, Louis: An outbreak of dermatitis from hair lacquer.....	1623
Sebrell, W. H.: The successful treatment of granulocytopenia and leukopenia in rats with crystalline folic acid.....	1542
Septic sore throat. (<i>See</i> United States—Monthly State reports.)	
Serum, human, jaundice following administration of [Oliphant, Gilliam, and Larson].....	1233
Sherrard, G. C.: The benefits accruing from the ratproof construction of vessels.....	1888
Shock and burns, experimental chemotherapy of. IV. Production of traumatic shock in mice. V. Therapy with mouse serum and sodium salts [Rosenthal].....	1429
Silicosis. (<i>See</i> United States—Monthly State reports.)	
Smallpox: (<i>See also</i> United States—Monthly State reports.)	
Foreign reports.....	1029, 1074, 1119, 1162, 1196, 1229, 1264, 1291, 1326, 1364, 1391, 1455, 1495, 1531, 1571, 1603, 1639, 1699, 1728, 1758, 1792, 1836, 1879, 1938
On a New Zealand hospital ship.....	1073
Weekly city reports.....	1025, 1069, 1115, 1157, 1191, 1225, 1259, 1287, 1320, 1359, 1387, 1419, 1450, 1490, 1526, 1566, 1597, 1633, 1661, 1695, 1723, 1753, 1788, 1831, 1874, 1902, 1933
Weekly State reports.....	1023, 1067, 1113, 1155, 1189, 1223, 1257, 1285, 1318, 1357, 1385, 1417, 1448, 1488, 1519, 1564, 1595, 1631, 1659, 1693, 1721, 1751, 1782, 1829, 1872, 1900, 1931
Smallpox, diphtheria, scarlet fever, and typhoid fever, extent of immunization and case histories for, in 200,000 surveyed families in 28 large cities [Collins and Council].....	1121
Smallpox in relation to State vaccination laws and regulations [Hampton].....	1771
Smith, Russell S.: Surveys of liquid wastes from munitions manufacturing—	
I. Trinitrotoluene (TNT) wastes.....	1365
II. Smokeless powder wastes.....	1373
III. Small arms ammunition.....	1393
IV. Tetryl wastes.....	1404
V. Nitroglycerine wastes.....	1409

	Page
Soap which indicates the presence of mercury fulminate [Mason and Botvinick].....	1183
Sodium fluoride, effect of topically applied, on dental caries experience [Knutson and Armstrong].....	1701
South Carolina. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
South Dakota. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Spain: Malaria.....	1074
Spicknall, Charles G.: The diet in Germany and the occupied countries during the second world war.....	1669
Spotted fevers of Brazil, Colombia, and the United States, experimental transmission of the, by the argasid tick <i>Ornithodoros nicolleti</i> [Davis].....	1742
Stead, F. M.: Study of an outbreak of food poisoning in a hospital in Galveston, Texas.....	1497
Steinhaus, Edward A.: <i>Salmonella enteritidis</i> : experimental transmission by the Rocky Mountain wood tick <i>Dermacentor andersoni</i> Stiles.....	1010
Stewart, Sarah E.: The mechanism of antitoxic immunity in <i>Clostridium perfringens</i> (Welchii) infections in guinea pigs.....	1277
Studies on trichinosis. XV. Summary of the findings of <i>Trichinella spiralis</i> in a random sampling and other samplings of the population of the United States [Wright, Kerr, and Jacobs].....	1293
Studies, statistical, a survey of, on the prevalence and incidence of mental disorder in sample populations [Lemkau, Tietze, and Cooper].....	1909
Sulfadiazine and atabrine, toxic effects of, in growing rats [Wright and Lillie].....	1242
Surveys of liquid wastes from munitions manufacturing:	
I. Trinitrotoluene (TNT) wastes [Smith and Walker].....	1365
II. Smokeless powder wastes [Smith and Walker].....	1373
III. Small arms ammunition [Smith and Walker].....	1393
IV. Tetryl wastes [Smith].....	1404
V. Nitroglycerine wastes [Smith].....	1409
Surveys of milk laboratories in war areas in the United States [Black]:	
I. Practices observed in making agar plate counts.....	1605
II. Practices observed in making direct microscopic examinations and methylene blue reduction tests.....	1641
III. Observations on sampling and health department practices relative to Lacteriological milk analysis.....	1681
Sweden: Notifiable diseases.....	1118, 1263, 1424, 1530, 1936
Switzerland: Notifiable diseases.....	1074, 1195, 1391, 1937

T

Tennessee. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Tetanus. (See United States—Monthly State reports.)	
Texas. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Texas, eastern, the occurrence of <i>Rickettsia diaporica</i> in <i>Amblyomma americanum</i> in. American Q fever: [Parker and Kohls].....	1510
Texas, Galveston, study of an outbreak of food poisoning in a hospital in [Lumsden, Nau, and Stead].....	1497
<i>Thrassis bacchi</i> (Roths.) and <i>Opisocrostitis bruneri</i> (Baker), report on the fleas, as vectors of plague [Prince].....	1013
Tick paralysis. (See United States—Monthly State reports.)	
Tietze, Christopher: A survey of statistical studies on the prevalence and incidence of mental disorder in sample populations.....	1909
Topping, Norman H.: A note on the rickettsioses in India.....	1208
Trachoma. (See United States—Monthly State reports.)	
<i>Trichinella spiralis</i> , summary of the findings of, in a random sampling and other samplings of the population of the United States. Studies on trichinosis. XV. [Wright, Kerr, and Jacobs].....	1293
Trichinosis: (See also United States—Monthly State reports.)	
Studies on. XV. Summary of the findings of <i>Trichinella spiralis</i> in a random sampling and other samplings of the population of the United States [Wright, Kerr, and Jacobs].....	1293

	Page
Trinitrotoluene poisoning, experimental, notes on the pathology of [Lillie]-----	1436
<i>Trypanosoma cruzi</i> , infection in monkeys with strains of, isolated in the United States [Davis]-----	1006
Tuberculosis:	
[All forms. (See United States—Monthly State reports.)	
Respiratory. (See United States—Monthly State reports.)	
Tuberculosis case finding, opportunities in the newer methods of [Hilleboe]-----	1094
Tuberculosis mortality in the United States, 1939-41 [Yerushalmy, Hilleboe, and Palmer]-----	1457
Tularemia: (See also United States—Monthly State reports.)	
Weekly city reports-----	1159,
1192, 1261, 1528, 1568, 1725, 1790, 1876, 1904, 1935	
Weekly State reports-----	1024, 1068, 1114, 1156, 1190, 1224,
1258, 1286, 1319, 1358, 1386, 1418, 1449, 1489, 1520, 1565, 1596,	
1632, 1660, 1694, 1722, 1752, 1783, 1830, 1873, 1901, 1932	
Tunisia: Infectious diseases—August 1943-----	1835
Typhoid and paratyphoid fever: (See also United States—Monthly State reports.)	
Weekly city reports-----	1025, 1069, 1115, 1157, 1191
1225, 1259, 1287, 1320, 1359, 1387, 1419, 1450, 1490, 1526, 1566,	
1597, 1633, 1661, 1695, 1723, 1753, 1788, 1831, 1874, 1902, 1933	
Weekly State reports-----	1023, 1067, 1113, 1155, 1189,
1223, 1257, 1285, 1318, 1357, 1385, 1417, 1448, 1488, 1519, 1564,	
1595, 1631, 1659, 1693, 1721, 1751, 1782, 1829, 1872, 1900, 1931	
Typhoid fever, diphtheria, smallpox, and scarlet fever, extent of immuniza- and case histories for, in 200,000 surveyed families in 28 large cities [Collins and Council]-----	1121
Typhus fever: (See also United States—Monthly State reports.)	
Foreign reports-----	1029, 1075, 1119, 1163,
1196, 1229, 1264, 1292, 1327, 1364, 1391, 1455, 1495, 1528, 1531,	
1571, 1603, 1639, 1665, 1699, 1728, 1759, 1836, 1879, 1907, 1939	
Weekly city reports-----	1027,
1071, 1117, 1159, 1192, 1226, 1261, 1289, 1322, 1360, 1452, 1568,	
1599, 1634, 1662, 1697, 1725, 1755, 1790, 1833, 1876, 1904, 1935	
Weekly State reports-----	1024, 1068, 1114, 1156, 1190,
1224, 1258, 1286, 1319, 1358, 1386, 1418, 1449, 1489, 1520, 1565,	
1596, 1632, 1660, 1694, 1722, 1752, 1783, 1830, 1873, 1901, 1932	

U

Undulant fever. (See United States—Monthly State reports.)	
United States:	
Communicable diseases—	
Current prevalence of-----	1061, 1215, 1482, 1625, 1745, 1891
Monthly State reports (published quarterly)-----	1521, 1784
Weekly city reports-----	1025, 1069, 1115, 1157, 1191,
1225, 1259, 1287, 1320, 1359, 1387, 1419, 1450, 1490, 1526, 1566,	
1597, 1633, 1661, 1695, 1723, 1753, 1788, 1831, 1874, 1902, 1933	
Weekly State reports-----	1021, 1065, 1110, 1153, 1187,
1221, 1255, 1283, 1315, 1355, 1383, 1415, 1446, 1486, 1517, 1562,	
1593, 1629, 1657, 1691, 1719, 1749, 1780, 1826, 1870, 1897, 1928	
Deaths—Large cities—Weekly reports (Weekly Mortality Index)---	1020,
1064, 1110, 1152, 1186, 1220, 1254, 1292, 1314, 1354, 1382, 1426,	
1439, 1485, 1516, 1561, 1592, 1636, 1666, 1690, 1718, 1748, 1778,	
1825, 1869, 1896, 1927	
(See also United States—Communicable diseases—Weekly city reports.)	
Health of the, at war-----	1281
Infection in monkeys with strains of <i>Trypanosoma cruzi</i> isolated in the [Davis]-----	1006
Influenza and pneumonia mortality in a group of 90 cities in the, August 1935-March 1943, with a summary for August 1920-March 1943 [Gover]-----	1033

United States—Continued.	Page
Mortality rates, provisional, for the first quarter of 1943.....	1439
Prevalence of poliomyelitis.....	1108, 1412
<i>Trichinella spiralis</i> in a random sampling and other samplings of the population of the, summary of the findings of. Studies on trichinosis. XV. [Wright, Kerr, and Jacobs].....	1293
Tuberculosis mortality in the, 1939-41 [Yerushalmy, Hilleboe, and Palmer].....	1457
United States, Colombia, and Brazil:	
Experimental transmission of the rickettsiae of the spotted fevers of, by the argasid tick <i>Ornithodoros nicolleti</i> [Davis].....	1742
Experimental transmission of the spotted fevers of the, by the argasid tick <i>Ornithodoros parkeri</i> [Davis].....	1201
The tick <i>Ornithodoros rudis</i> as a host to the rickettsiae of the spotted fevers of [Davis].....	1016
Utah. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
V	
Vermont. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Vincent's infection. (See United States—Monthly State reports.)	
Virgin Islands of the United States: Notifiable diseases.....	1227, 1635
Virginia. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Vital statistics. (See Name of country.)	
W	
Walker, W. W.: Surveys of liquid wastes from munitions manufacturing—	
I. Trinitrotoluene (TNT) wastes.....	1365
II. Smokeless powder wastes.....	1373
III. Small arms ammunition.....	1393
War and the distribution of physicians [Perrott and Davis].....	1545
War, health of the United States at.....	1281
Washington. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Water contaminated with chemical warfare agents, arsenic in, the detection and analysis of [Ruchhoft, Placak, and Schott].....	1761
Wattie, Elsie: Influence of pH and temperature on the survival of coliforms and enteric pathogens when exposed to free chlorine.....	1837
Weil's disease. (See United States—Monthly State reports.)	
West Virginia. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Whooping cough: (See also United States—Monthly State reports.)	
Weekly city reports.....	1025, 1069, 1115, 1157, 1191, 1225, 1259, 1287, 1320, 1359, 1387, 1419, 1450, 1490, 1526, 1566, 1597, 1633, 1661, 1695, 1723, 1753, 1788, 1831, 1874, 1902, 1933
Weekly State reports.....	1024, 1068, 1114, 1156, 1190, 1224, 1258, 1286, 1319, 1358, 1386, 1418, 1449, 1489, 1520, 1565, 1596, 1632, 1660, 1694, 1722, 1752, 1783, 1830, 1873, 1901, 1932
Wisconsin. (See Name of disease—Weekly State reports; United States—Monthly State reports.)	
Wright, C. I.: Toxic effects of atabrine and sulfadiazine in growing rats..	1242
Wright, Willard H.: Studies on trichinosis. XV. Summary of the findings of <i>Trichinella spiralis</i> in a random sampling and other samplings of the population of the United States.....	1293
Wyoming: (See also Name of disease—Weekly State reports; United States—Monthly State reports.)	
Plague infection in.....	1290, 1600

Y

Yellow fever: Foreign reports.....	1119, 1163, 1197, 1229, 1327, 1364, 1392, 1425, 1456, 1571, 1603, 1640, 1665, 1699, 1728, 1760, 1836, 1880, 1940
Yerushalmy, J.: Tuberculosis mortality in the United States, 1939-41....	1457
Young, Martin D.: Carbarsone treatment for <i>Balantidium coli</i> infections..	1272

ge
39
12

93

57

42

01

16

35

5
3
3
5
1

1

7

7
3
6
2

2

3

0

7
7
4

